

SN 2007uy – metamorphosis of an aspheric Type Ib explosion

Rupak Roy¹*, Brijesh Kumar¹, Justyn R. Maund²†, Patricia Schady³, Felipe Olivares E.³, Daniele Malesani⁴, Giorgos Leloudas^{4,5}, Sumana Nandi¹, Nial Tanvir⁶, Dan Milisavljevic⁷, Jens Hjorth⁴, Kuntal Misra¹, Brajesh Kumar¹, S. B. Pandey¹, Ram Sagar¹, H. C. Chandola⁸

¹*Aryabhatta Research Institute of Observational Sciences, Manora Peak, Nainital, 263 002, India*

²*Astrophysics Research Centre, School of Mathematics and Physics, Queens University Belfast, Belfast BT7 1NN, UK*

³*Max-Planck-Institute für extraterrestrische Physik, Giessenbachstraße 1, 85748 Garching, Germany*

⁴*Dark Cosmology Centre, Niels Bohr Institute, University of Copenhagen, Juliane Maries vej 30, 2100 Copenhagen Ø, Denmark*

⁵*The Oskar Klein Centre, Department of Physics, Stockholm University, Albanova University Centre, 10691, Stockholm, Sweden.*

⁶*Department of Physics and Astronomy, University of Leicester, Leicester, LE1 7RH, United Kingdom*

⁷*Department of Physics and Astronomy, Dartmouth College, Hanover, NH 03755*

⁸*Department of Physics, Kumaun University, Nainital, India*

Accepted ??; Received ??

ABSTRACT

The supernovae (SNe) of Type Ibc are rare and the detailed characteristics of these explosions have been studied only for a few events. Unlike Type II SNe, the progenitors of Type Ibc have never been detected in pre-explosion images. So, to understand the nature of their progenitors and the characteristics of the explosions, investigation of proximate events are necessary. Here we present the results of multi-wavelength observations of Type Ib SN 2007uy in the nearby (~ 29.5 Mpc) galaxy NGC 2770. Analysis of the photometric observations revealed this explosion as an energetic event with peak absolute R band magnitude -18.5 ± 0.16 , which is about one mag brighter than the mean value (-17.6 ± 0.6) derived for well observed Type Ibc events. The SN is highly extinguished, $E(B - V) = 0.63 \pm 0.15$ mag, mainly due to foreground material present in the host galaxy. From optical light curve modeling we determine that about $0.3 M_{\odot}$ radioactive ^{56}Ni is produced and roughly $4.4 M_{\odot}$ material is ejected during this explosion with liberated energy $\sim 15 \times 10^{51}$ erg, indicating the event to be an energetic one. Through optical spectroscopy, we have noticed a clear aspheric evolution of several line forming regions, but no dependency of asymmetry is seen on the distribution of ^{56}Ni inside the ejecta. The SN shock interaction with the circumburst material (CSM) is clearly noticeable in radio follow-up, presenting a Synchrotron Self Absorption (SSA) dominated light curve with a contribution of Free Free Absorption (FFA) during the early phases. Assuming a Wolf-Rayet (WR) star, with wind velocity $\gtrsim 10^3 \text{ km s}^{-1}$, as a progenitor, we derive a lower limit to the mass loss rate inferred from the radio data as $\dot{M} \gtrsim 2.4 \times 10^{-5} M_{\odot} \text{ yr}^{-1}$, which is consistent with the results obtained for other Type Ibc SNe bright at radio frequencies.

Key words: supernovae: general – supernovae: individual: SN 2007uy – galaxies: individual: NGC 2770

1 INTRODUCTION

Core-collapse supernovae (CCSNe) mark the end stage of the evolution of massive stars having initial masses greater than $8 M_{\odot}$ (Heger et al. 2003; Burrows 2013). Stars, more

* e-mail: roy@aries.res.in, rupakroy1980@gmail.com

† Royal Society Research Fellow

massive than $10M_{\odot}$ produce an Fe-core, and then expel their outer layers in a catastrophic explosion (e.g., Pumo et al. 2009). Type Ib SNe are a class of CCSNe which are spectroscopically defined by the absence of hydrogen (H) and presence of helium (He) in their early phase optical spectra. It is generally accepted that Type Ib SNe along with Type Ic (absence of H and He) are formed from evolved high mass progenitors like Wolf-Rayet (WR) stars which have liberated their outer shells through pre-SN winds. The outer shells of H & He may also be stripped through mass transfer due to Roche-lobe overflow in a binary system. Type Ibc SNe are commonly referred to as “Stripped-envelope Supernovae” (Clocchiatti & Wheeler 1997). The progenitors of Type Ibc SNe have not been observed directly in pre-explosion high resolution images. There are only a few cases, like SNe 2000ds, 2000ew, 2001B and 2004gt where upper-limits on the flux of the Type Ibc progenitors have been reported (Maund & Smartt 2005; Maund et al. 2005; Eldridge et al. 2013). After analyzing the evolutionary models of massive He stars and comparing the results with Galactic WR stars, it has been found that during pre-SN stage, progenitors of Type Ibc SNe have surface properties that resemble those of hot Galactic WR stars of WO sub-type, which are visually faint, $M_V \approx -1.5$ to -2.5 mag, despite of a high bolometric luminosity, $L/L_{\odot} = 5.6 - 5.7$ (Yoon et al. 2012). Detection of Type Ic progenitors is even more challenging than that of Type Ib. Monitoring of this class of SNe is crucial, not only to investigate the post-explosion phenomenon, but also to understand the pre-explosion properties of the progenitors.

With a volumetric sample of nearby CCSNe discovered by the Lick Observatory Supernova Search (LOSS) program, Smith et al. (2011) found that only about 7% of total nearby CCSNe are of Type Ib. Furthermore they argued that most of the Type Ib are formed through mass transfer in a binary system rather than through the progenitor’s wind and that the true progenitors of SNe Ib must extend to a much lower range of initial masses than classical WR stars.

Apart from the issues related to progenitors and the pre-explosion properties, there are many unresolved issues concerning the post-explosion properties. One of them is the explosion geometry. Spectropolarimetry has revealed CCSNe to be more aspheric in their inner layers with a preferred direction of explosion in comparison with thermonuclear events, which instead show asphericity in their outer shells (Wang & Wheeler 2008). Among CCSNe, Type IIP events show a moderate non-zero polarization in the plateau phase, whereas Type Ibc SNe are aspheric explosions with substantially higher intrinsic polarization from the beginning (Wang et al. 2001; Maund et al. 2007). The asphericity is explained as being due to clumpy structure or jet-like propagation of the ionizing radioactive Ni and Co source, concentrated at the inner portion of the spherically symmetric ejecta (Chugai 1992; Höflich et al. 2001; Maund et al. 2007). In the light of mounting evidence of the asphericity of Type Ibc events, the double horned nature of the [O I] $\lambda\lambda 6300, 6364$ profile was explained as a jet/torus like structure (Mazzali et al. 2005; Modjaz et al. 2008; Maeda et al. 2008). Though, alternatively, it can be explained as a combined effect of two [O I] distributions – (i) a central, symmetric distribution of [O I]-rich ejecta, (ii) a clumpy or shell of [O I]-rich material traveling in the front facing hemisphere (although this model also has some drawbacks; see




Figure 1. Identification chart of SN 2007uy. The image is about 10 arcmin on a side taken in V-band with the 104-cm Sampurnanand telescope at ARIES, Nainital. The local second-order stars are numbered. North is up and east is to the left.

Milisavljevic et al. 2010; Taubenberger et al. 2009). Clearly, individual case studies are important to understand the preferred mechanism behind the ambiguous behavior of [O I] lines.

SN 2007uy, a Type Ib event, was first discovered by Yoji Hirose on 2007, December 31.7 UT at an unfiltered magnitude 17.2 (Nakano et al. 2008) in the nearby galaxy NGC 2770 ($z \sim 0.007$, distance ~ 29.5 Mpc). There are numerous intervening host galaxy H II regions along the direction of the SN (Gorosabel et al. 2011) and no object was found at the transient location in the red band DSS image observed between 1998 and 2000. Blondin & Calkins (2008) reported the first optical spectroscopic observation of this event taken on 2008, January 3.40 UT, classifying this transient as a type Ib event similar to SN 2004gq. The discovery date corresponds to JD = 2454466.17. According to initial spectroscopy (Blondin & Calkins 2008) the explosion happened within 7 days prior to its discovery. Modeling of radio data (§7) also indicates that the event was discovered within 4 days of the explosion. Here we adopt a conservative value of the explosion epoch to be 4 days prior the detection, i.e. JD = 2454462.17. In rest of the work, if not mentioned, all the phases will be quoted with respect to this epoch. The field of SN 2007uy is presented in figure 1 and the properties of the host galaxy and the SN are provided in Table 1. A detailed description of the host can be found elsewhere (Thöne et al. 2009). SN 2007uy was also detected in radio bands at +10d (Soderberg 2008) with an X-band (8.46 GHz) flux density of 0.29 ± 0.03 mJy. The SN was also detected at X-ray wavelengths (Pooley & Soderberg 2008).

In this paper, we present photometric follow-up of SN 2007uy in near-ultraviolet (NUV), optical and near-infrared (NIR) bands and low resolution spectroscopic follow-up observations in optical. We summarize the observations and data reduction procedure in §2. We study the spectroscopic evolution in §3. The distance and reddening have been estimated in section §4. In §5 photometric evolution is investi-

Table 1. Properties of the host galaxy NGC 2770 and SN 2007uy.

Parameters	Value	Ref. ^a
NGC 2770:		
Type	SABc	1
Position	$\alpha_{J2000} = 09^h09^m33^s.68$ $\delta_{J2000} = +33^\circ07'24''.7$	1
Abs. magnitude	$M_B = -20.78$ mag	1
Distance	$D = 29.5 \pm 1.8$ Mpc	§4
Scale	$1'' \sim 136$ pc, $1' \sim 8.2$ kpc	
Distance modulus	$\mu \sim 32.4$ mag	
Apparent radius	$r_{25} = 1.73$ (~ 15.09 kpc)	1
Inclination angle	$\Theta_{inc} = 82.3^\circ$	1
Position angle	$\Theta_{maj} = 146.1^\circ$	1
Heliocentric velocity	$cz_{helio} = 1947 \pm 2$ km s ⁻¹	1
SN 2007uy:		
Position	$\alpha_{J2000} = 09^h09^m35^s.40$ $\delta_{J2000} = +33^\circ07'09''.9$	2
Location	20'6 E, 15'5 S	2
Deprojected radius	$r_{SN} = 42''.44$ (~ 5.3 kpc)	§4
Discovery date (UT)	31.7 December 2007 (JD 2454466.17)	2
Explosion epoch:	~ 4 days prior to discovery (JD 2454462.17)	§1
Total reddening :	$E(B - V) = 0.63 \pm 0.15$ mag	§4

^a(1) HyperLEDA - <http://leda.univ-lyon1.fr>; (2) Nakano et al. (2008).

gated, while the section §6 presents the evolution of colour and bolometric light. The main physical parameters of the explosion and the characteristics of the progenitor are derived and discussed in §7 and §8.

2 OBSERVATIONS AND DATA REDUCTION

2.1 Near-ultraviolet, optical and near-infrared multiband photometry

The Ultraviolet Optical Telescope (UVOT) monitored the SN at 30 phases between +9d and +75d. The observations were carried out in *uvw2* ($\lambda_c = 2030\text{\AA}$), *uvm2* ($\lambda_c = 2231\text{\AA}$), *uvw1* ($\lambda_c = 2634\text{\AA}$), *u* ($\lambda_c = 3501\text{\AA}$), *b* ($\lambda_c = 4329\text{\AA}$) and *v* ($\lambda_c = 5402\text{\AA}$) bands (see table 2 for details). We obtained the UVOT data from the *Swift* Data Archive. The photometry was done using standard procedures using HEASOFT routines¹ and it was calibrated to the UVOT

¹ The *Swift* data have had bad pixels identified, mod-8 noise corrected, and have been transformed into FK5 coordinates. We used the standard UVOT data analysis software distributed with HEASOFT 6.10 along with the standard calibration data. As long as the source had a count rate greater than 0.5 cts s^{-1} , photometry was done using ‘uvotsource’ with a standard circular aperture of radius $5''$ and a circular background region with a radius of $15''$. Below this threshold, a $3''.5$ radius was used and an aperture correction was applied. The background region was selected to

photometric system following the procedure described in Poole et al. (2008). To remove the contribution of the underlying host galaxy, we measured the host galaxy flux at the position of the SN from late time UVOT observations, with no contribution from the SN. This additional flux was then subtracted from SN photometric measurements. For all observations the source was close to the centre of the field-of-view, and differences in the PSF (Point Spread Function) between observations were, therefore, negligible.

The optical broadband Johnson *UBV* and Cousins *RI* follow-up observations were performed at 27 phases between $\sim +16\text{d}$ to $+130\text{d}$ using the 104-cm Sampurnanand Telescope (ST)+imaging camera² at Nainital, India and the 2.56 m Nordic Optical Telescope (NOT) + Andalucia Faint Object Spectrograph and Camera (ALFOSC) at La Palma, Spain. Several exposures, in the range 100s to 300s, were taken and in order to increase the signal-to-noise-ratio (SNR), photometry was performed on the co-added frames. The pre-processing of raw data were performed through standard data reduction software *IRAF*³ and photometry was performed using the stand-alone version of *DAOPHOT*⁴ (Stetson 1987).

The SN was surrounded by a star forming knot and the host galaxy is highly inclined ($\sim 82.3^\circ$) with respect to the line of sight. The SN flux was, therefore highly contaminated by the background. In order to account for this, late-time (~ 36 months) high SNR images of the host galaxy in *UBVRI* bands, obtained from the NOT under good photometric conditions, were used to map the galaxy flux, and a template subtraction technique (Roy et al. 2011a,b) was adopted to remove the galaxy contributions. Before flux subtraction, both NOT and ST images were brought to a common ‘plate-scale’ using the ‘magnify’ task provided in the *IRAF* package. The instrumental magnitudes of the SN were derived from the template subtracted SN frames using the PSF fitting method.

The field of SN 2007uy was calibrated from ST in the *BVRI* bands using Landolt (1992) standard stars of the fields of PG1633+009 and PG1047+003 observed on the night of March 02, 2008 under moderate seeing (FWHM $\sim 2''$ in R band) and transparent sky conditions. We used mean values of the atmospheric extinction coefficients of the site (namely 0.28, 0.17, 0.11 and 0.07 mag per unit airmass for the *B*, *V*, *R* and *I* bands, from Kumar et al. 2000) with typical standard deviations between the transformed and the standard magnitudes of Landolt stars of 0.04 in *B*, 0.02

have similar background properties to those at the location of the SN, and to be free of contaminating sources.

² A 2048×2048 CCD camera having $24 \times 24 \mu\text{m}$ chip size and with a plate scale $0''.38/\text{pixel}$, mounted at the f/13 Cassegrain focus of the telescope was used for observation. The gain and readout noise of the CCD camera are 10 electrons per analog-to-digital unit and 5.3 electrons respectively. To improve the signal-to-noise ratio, we performed the observations in a binned mode of 2×2 pixel.

³ *IRAF* stands for Image Reduction and Analysis Facility distributed by the National Optical Astronomy Observatories which is operated by the Association of Universities for research in Astronomy, Inc. under co-operative agreement with the National Science Foundation

⁴ *DAOPHOT* stands for Dominion Astrophysical Observatory Photometry

Table 2. *swift*/UVOT photometry of SN 2007uy.

UT Date (yy/mm/dd)	JD 2454000+	Phase ^a (day)	uvw2 (mag)	uvm2 (mag)	uvw1 (mag)	u (mag)	b (mag)	v (mag)
2008/01/06.15	471.65	+9	21.45	24.33	18.61±0.18	16.89±0.06	17.28±0.06	16.62±0.08
2008/01/06.49	471.99	+10	20.29±0.39	21.31±4.25	18.45±0.10	16.93±0.04	17.02±0.03	16.47±0.04
2008/01/09.71	475.21	+13	20.05±0.35	20.15±0.58	18.23±0.10	16.78±0.04	16.81±0.03	16.09±0.04
2008/01/11.38	476.88	+14	20.33±0.27	20.69±0.65	18.37±0.06	16.69±0.02	16.79±0.02	15.96±0.02
2008/01/11.99	477.49	+15	21.28±0.82	>20.57	18.51±0.07	16.72±0.02	16.81±0.02	15.93±0.02
2008/01/12.57	478.07	+15	20.48±0.40	21.40±3.72	18.45±0.09	16.76±0.03	16.77±0.02	15.85±0.03
2008/01/13.21	478.71	+16	20.59±0.47	22.63	18.50±0.09	16.76±0.03	16.80±0.03	15.93±0.03
2008/01/13.89	479.39	+17	20.34±0.29	20.81±0.68	18.41±0.07	16.78±0.03	16.81±0.02	15.81±0.02
2008/01/14.73	480.23	+18	21.41±0.87	>20.59	18.61±0.08	16.88±0.03	16.77±0.02	15.80±0.02
2008/01/15.49	480.99	+18	21.32±1.00	>20.36	18.95±0.12	17.03±0.04	16.81±0.02	15.79±0.02
2008/01/16.29	481.79	+19	20.77±0.47	>20.36	18.85±0.11	17.04±0.04	16.81±0.02	15.78±0.02
2008/01/17.14	482.64	+20	21.43	>19.88	18.93±0.09	17.15±0.03	16.90±0.03	15.77±0.02
2008/01/17.81	483.31	+21	21.53±0.96	22.97	18.94±0.08	17.27±0.03	16.89±0.02	15.73±0.02
2008/01/18.85	484.35	+22	>20.87	22.19	—	—	—	—
2008/01/21.12	486.62	+23	22.92	24.98	19.62±0.11	18.07±0.05	17.26±0.03	15.89±0.02
2008/01/21.76	487.26	+25	—	—	19.71±0.40	18.30±0.17	17.25±0.07	15.94±0.07
2008/01/25.17	490.67	+28	—	—	>20.64	—	—	—
2008/01/26.41	491.91	+29	—	—	20.84±0.29	19.33±0.10	17.88±0.03	16.28±0.02
2008/01/27.98	493.48	+31	—	—	>20.91	—	—	—
2008/01/29.35	494.85	+32	—	—	>20.78	—	—	—
2008/01/30.81	496.31	+35	—	—	22.04±0.50	20.78±0.58	18.51±0.06	16.66±0.03
2008/02/01.93	498.43	+37	—	—	>20.87	21.59±1.57	18.53±0.06	16.86±0.03
2008/02/03.93	500.43	+39	—	—	19.90±1.25	—	—	—
2008/02/08.95	505.45	+43	—	—	22.39±0.50	—	—	—
2008/02/10.89	507.39	+45	—	—	—	—	19.05±0.10	—
2008/02/13.03	509.53	+47	—	—	—	—	18.95±0.09	17.27±0.05
2008/02/16.11	512.61	+50	—	—	—	—	—	—
2008/02/22.30	518.80	+57	—	—	—	—	—	—
2008/02/25.79	522.29	+60	—	—	—	—	18.96±1.04	17.72±0.09
2008/03/12.60	538.10	+75	—	—	—	—	—	17.84±0.12

^aWith reference to the epoch of explosion JD 2454462.17**Table 3.** Photometry of local standard stars in the field of SN 2007uy^a

Star ID	α_{2000} (h m s)	δ_{2000} (deg m s)	B (mag)	V (mag)	R (mag)	I (mag)
1	09:09:42.48	33:05:07.1	16.37±0.04	15.56±0.02	15.06±0.02	14.64±0.01
2	09:09:44.25	33:05:11.8	17.64±0.04	16.98±0.02	16.62±0.03	16.27±0.01
3	09:09:37.26	33:09:34.9	17.85±0.04	17.34±0.03	17.01±0.03	16.69±0.02
4	09:09:36.84	33:05:46.0	18.00±0.05	17.26±0.02	16.84±0.03	16.43±0.01
5	09:09:36.12	33:05:42.9	18.11±0.05	17.48±0.03	17.13±0.03	16.81±0.02
6	09:09:31.93	33:05:19.5	19.01±0.05	17.93±0.04	17.26±0.04	16.74±0.01
7	09:09:18.95	33:08:33.4	17.54±0.04	17.15±0.02	16.87±0.03	16.57±0.01
8	09:09:14.29	33:07:16.0	18.85±0.05	17.49±0.03	16.63±0.03	15.92±0.01
9	09:09:11.82	33:06:34.6	19.09±0.05	18.02±0.04	17.36±0.04	16.86±0.01
10	09:09:10.49	33:06:25.6	19.83±0.05	18.72±0.04	17.97±0.04	17.37±0.02

^aThe U band data is standardized with respect to the field standards mentioned in Malesani et al. (2009). Errors are 1σ uncertainties and it include both photometric as well as calibration errors.

in V and R and 0.01 in I band. A sample of 10 bright and isolated non-variable stars in the field of SN 2007uy were used as local standards to derive the zero points for the SN at each epoch. The location and magnitudes of these local standards are listed in Table 3. These secondary standards are marked in figure 1. For U band, the calibration of Malesani et al. (2009) is used. Table 4 lists the final pho-

tometry of SN 2007uy in the $UBVRI$ bands at 37 phases between +16d and +129d.

The field was also monitored in JHK NIR bands at 8 phases between +16d and +87d using United Kingdom Infrared Telescope (UKIRT) with Wide-Field Camera (WFCAM) as a backend detector (Hirst et al. 2006). The dithered images were processed with standard tasks in *IRAF* and photometry was performed with *DAOPHOT*

Table 4. *UBVRI* photometry of SN 2007uy.

UT Date (yyyy/mm/dd)	JD 2454000+	Phase ^a (day)	<i>U</i> (mag)	<i>B</i> (mag)	<i>V</i> (mag)	<i>R</i> (mag)	<i>I</i> (mag)	Telescope ^b
2008/01/12.94	478.44	+16	16.64±0.01	16.78±0.01	15.79±0.01	15.54±0.01	—	NOT
2008/01/13.99	479.71	+17	—	16.77±0.02	15.81±0.03	15.51±0.03	15.53±0.03	ST
2008/01/14.85	480.58	+18	—	16.74±0.03	15.78±0.03	15.46±0.02	15.43±0.04	ST
2008/01/15.19	480.69	+18	16.69±0.03	16.71±0.02	15.76±0.01	15.38±0.02	15.31±0.01	NOT
2008/01/15.99	481.72	+19	—	16.78±0.02	15.74±0.02	15.44±0.01	15.36±0.03	ST
2008/01/17.19	482.69	+20	16.95±0.01	16.80±0.01	15.81±0.01	15.40±0.01	15.21±0.01	NOT
2008/01/28.94	494.44	+32	—	17.98±0.01	16.45±0.01	15.72±0.01	15.50±0.01	NOT
2008/01/29.84	495.57	+33	—	18.19±0.02	16.44±0.02	15.81±0.02	15.59±0.02	ST
2008/01/30.19	495.69	+33	—	—	16.53±0.02	15.79±0.01	15.58±0.01	NOT
2008/01/31.94	497.44	+35	19.08±0.02	18.22±0.01	16.60±0.01	15.89±0.01	15.62±0.02	NOT
2008/01/31.87	497.60	+35	—	18.28±0.03	16.54±0.03	15.91±0.03	15.62±0.04	ST
2008/02/01.94	498.44	+36	19.20±0.03	18.29±0.02	16.66±0.02	15.96±0.02	15.66±0.02	NOT
2008/02/03.94	500.44	+38	19.32±0.03	18.41±0.03	16.86±0.02	16.08±0.02	15.77±0.02	NOT
2008/02/04.88	501.61	+39	—	18.50±0.06	16.82±0.02	16.11±0.02	15.83±0.03	ST
2008/02/09.75	506.48	+44	—	18.70±0.03	17.07±0.03	16.40±0.03	15.97±0.04	ST
2008/02/05.86	512.59	+50	—	18.84±0.03	17.33±0.01	—	16.18±0.03	ST
2008/02/18.94	515.44	+53	19.75±0.04	19.01±0.03	17.33±0.03	16.66±0.03	16.26±0.03	NOT
2008/02/23.65	520.37	+57	—	19.27±0.05	17.64±0.04	16.84±0.03	16.53±0.07	ST
2008/02/24.62	521.35	+58	—	—	17.52±0.02	16.88±0.02	16.44±0.03	ST
2008/02/29.94	526.44	+64	—	—	17.58±0.03	—	16.48±0.03	NOT
2008/03/02.75	529.45	+66	—	19.22±0.07	—	—	—	ST
2008/03/03.72	531.53	+67	—	19.28±0.06	17.66±0.04	17.05±0.03	16.54±0.05	ST
2008/03/05.80	533.61	+69	—	19.29±0.03	—	17.05±0.04	16.53±0.06	ST
2008/03/17.69	543.19	+81	—	19.31±0.04	17.81±0.03	17.29±0.03	16.76±0.03	NOT
2008/03/05.80	587.36	+124	—	—	18.67±0.02	18.32±0.03	18.02±0.06	ST
2008/05/02.61	589.34	+126	—	—	—	18.51±0.05	—	ST
2008/05/05.62	592.35	+129	—	—	18.84±0.03	—	—	ST

^aWith reference to the epoch of explosion JD 2454462.17.^bThe photometric observations are taken with the 1-m Sampurnanand Telescope (ST), ARIES, Nainital and 2.6-m Nordic Optical Telescope (NOT) with ALFOSC detector. Errors in magnitude denote 1σ uncertainty.**Table 5.** Near-infrared *JHK* photometry of SN 2007uy.

UT Date (yyyy/mm/dd)	JD 2454000+	Phase ^a (day)	<i>J</i> (mag)	<i>H</i> (mag)	<i>K</i> (mag)
2008/01/12.67	478.17	+16	14.48±0.25	14.98±0.25	14.98±0.17
2008/01/14.68	480.18	+18	14.32±0.19	14.80±0.24	14.71±0.14
2008/01/15.66	481.16	+19	14.26±0.19	14.73±0.24	14.67±0.15
2008/01/17.69	483.19	+21	14.12±0.32	14.64±0.28	14.51±0.22
2008/01/23.67	489.17	+27	14.13±0.25	14.60±0.25	14.51±0.25
2008/02/15.92	512.42	+50	15.76±0.07	15.38±0.02	15.06±0.18
2008/02/24.92	521.42	+59	16.07±0.06	15.58±0.03	15.32±0.19
2008/03/23.92	549.42	+87	17.03±0.07	16.36±0.03	16.21±0.19

^aWith reference to the epoch of explosion JD 2454462.17.

routines. The field was calibrated with respect to nearby 2MASS stars, and the calibrated *JHK* magnitudes of SN 2007uy is given in Table 5.

2.2 Radio Data

The transient was first detected in radio on 2008 January 6.18 UT, using the Very Large Array (VLA) at 8.46 GHz, with a flux density of $290 \pm 30 \mu\text{Jy}$ (Soderberg 2008). Although the VLA observations originally targeted SN 2007uy, after January 10.2 UT the field of observation was centred on another bright transient, SN 2008D, discovered in the

same galaxy. Due to larger primary beam size of VLA in L (30'), C (9') and X (5.4') bands, the locations of both transients were within VLA field of view. Here, L band corresponds to 1.34-1.73 GHz, C band corresponds to 4.5-5.0 GHz and X band corresponds to 8.0-8.8 GHz radiation. We used the NRAO archival facility to fetch these data sets and reduced the data using standard routines of 'Astronomical Image Processing System' (AIPS)⁵. The transient was

⁵ Astronomical Image Processing System (AIPS) has been developed by the National Radio Astronomical Observatories (NRAO), USA

Table 6. Log of radio observation of SN 2007uy from VLA in 4.8 & 8.4 GHz.

UT Date (yy/mm/dd)	JD 2454000+	Phase ^a (day)	Frequency (GHz)	Flux (mJy)	Flux error ^b (mJy)
2008/01/06	471.76	+09	8.4	0.362	0.044
2008/01/07	473.11	+10	8.4	0.306	0.070
2008/01/11	476.82	+14	8.4	0.459	0.469
2008/01/14	479.76	+17	8.4	0.513	0.040
2008/01/16	481.80	+19	4.8	0.515	0.050
2008/01/16	481.83	+19	8.4	0.688	0.057
2008/01/17	482.90	+20	4.8	0.403	0.037
2008/01/17	482.92	+20	8.4	0.566	0.053
2008/01/20	485.66	+23	4.8	0.562	0.044
2008/01/21	486.66	+24	8.4	0.724	0.060
2008/01/21	486.67	+24	4.8	0.799	0.055
2008/01/23	488.66	+26	4.8	0.770	0.060
2008/01/23	488.67	+26	8.4	0.721	0.076
2008/01/25	490.67	+28	8.4	0.797	0.056
2008/01/25	490.68	+28	4.8	0.875	0.068
2008/01/27	492.92	+30	8.4	0.965	0.081
2008/01/27	492.93	+30	4.8	0.821	0.050
2008/01/30	495.82	+33	8.4	0.953	0.070
2008/02/01	497.73	+35	4.8	1.216	0.085
2008/02/01	497.74	+35	8.4	0.968	0.068
2008/02/03	499.70	+37	8.4	1.044	0.075
2008/02/03	499.71	+37	4.8	1.072	0.065
2008/02/08	504.72	+42	4.8	1.263	0.078
2008/02/08	504.73	+42	8.4	1.011	0.075
2008/02/14	510.79	+48	8.4	0.872	0.063
2008/02/14	510.80	+48	4.8	1.249	0.085
2008/02/21	517.92	+55	4.8	1.276	0.102
2008/02/21	517.93	+55	8.4	0.522	0.048
2008/02/24	520.65	+58	4.8	1.432	0.082
2008/02/24	520.66	+58	8.4	0.677	0.046
2008/03/07	532.60	+70	4.8	1.339	0.106
2008/03/07	532.62	+70	8.4	0.596	0.062
2008/03/21	546.58	+84	4.8	1.103	0.089
2008/03/22	547.65	+85	8.4	0.351	0.049
2008/04/14	570.53	+108	4.8	0.814	0.120
2008/05/04	591.36	+129	4.8	0.457	0.107

^aWith reference to the explosion epoch JD 2454466.17

^bThe Flux errors are measured using the expression $\sigma_f^2 = (\epsilon.S_0)^2 + \sigma_0^2 + \sigma_{S_0}^2$. Here S_0 is the observed flux density, σ_0 is the RMS noise of the radio sky, σ_{S_0} is the error associated with σ_0 and ϵ is the fraction that accounts the error in VLA flux calibration. For 4.8 and 8.4 GHz observations value of ϵ is 0.05. For the epochs where measured flux density is less than 3 times of the corresponding value of σ_f , we have considered the flux density as the upper limit for our measurement.

not detected in the radio U (14.4-15.4 GHz) band, but was prominent in the C and X band images. The new VLA data set is presented in Table 6. For further analysis we have used the literature data (van der Horst et al. 2011) along with the VLA archival data.

2.3 Low-resolution optical spectroscopy

The spectroscopic observations of SN 2007uy were carried out at 7 epochs between +17d to +392d. The spectral data for +17d were acquired with NOT/ALFOSC on January 13, 2008. The spectra for the epochs +32d, +58d, +122d & +392d are based on the archival data obtained through ‘ESO Science Archive Portal’ which were acquired using 8m VLT and 3.6m NTT. The spectral data for +96d and +162d were taken from Milisavljevic et al. (2010). These were primarily acquired on 1st April and 6th June, 2008 using the ‘MMT-

BLUECHANNEL’ detector at MMT. A journal of spectroscopic observations is presented in Table 7.

All the raw optical data were processed using the standard tasks in *IRAF*. Bias and flat-fielding were performed on each frame. Cosmic ray rejection on each frame was done by using Laplacian kernel detection routine LACOSMIC (van Dokkum 2001)⁶. Images were co-added to improve the signal-to-noise ratio and one-dimensional spectra were extracted from co-added frames using the *apall* task in *IRAF* (Horne 1986). Wavelength calibration was performed using the *identify* task and the fifth order fits were used to achieve a typical RMS scatter of 0.1Å (i.e., 60 km s⁻¹ at 5000Å). The position of the O I emission skyline at 5577Å was used to check the wavelength calibration and deviations were found

⁶ <http://www.astro.yale.edu/dokkum/lacosmic>




Figure 2. Spectroscopic evolution of SN 2007uy. All the spectra have been normalized with respect to the peak flux of the underlying $H\alpha$ feature and a constant offset has been applied to present them clearly. The +162d and +392d spectra have been multiplied by a factor of 2 to enlarge several tiny features. The dotted vertical line represents the position of $H\alpha$ and confirms the wavelength calibration within the limits of the spectral resolution.

to lie between 0.5 to $1\text{\AA}$ and this was corrected by applying a linear shift in wavelength.

3 SPECTROSCOPIC EVOLUTION

In order to proceed further with the spectral analysis, the wavelength calibrated spectra were corrected for the recession velocity of the host galaxy using the prominent Balmer emission lines arising from the underlying HII regions. In this way both the redshifts due to recession and rotation of the galaxy have been accounted for. All the spectra were

normalized with respect to the peak flux of the underlying $H\alpha$ feature and a constant offset was applied to present them clearly. In figure 2, the spectra of SN 2007uy is presented at 7 epochs spanning between +17d and +392d. The +17d spectrum was taken near maximum light, while +162d and +392d are late nebular phase spectra. The latter are enlarged by a factor of 2 for clarity. Spectral features are mainly identified as per previously published line identification lists for Type Ibc SNe (Cox 2000 and references therein), though high degree of line-blending and line-blanking limit the detections. The dotted vertical line rep-

Table 7. Journal of spectroscopic observations of SN 2007uy ^a.

UT Date (yy/mm/dd)	Phase ^b (days)	Range μm	Telescope ^c	Grating (gr mm ⁻¹)	Slit width ($''$)	Dispersion ($\text{\AA pix}^{-1}$)	Exposure (s)	SNR ^d (pix ⁻¹)
2008/01/13	+17	0.32–0.91	NOT	300	1.3	3.0	1200	135
2008/01/28	+32	0.38–0.92	NTT	300	1.0	1.7	1200	170
2008/02/22	+58	0.45–0.87	VLT	300	1.3	1.7	900	225
2008/04/01	+96	0.32–0.80	MMT	300	1.0	2.0	600×4	110
2008/04/26	+122	0.35–0.87	VLT	300	1.3	1.7	2940	250
2008/06/06	+162	0.38–0.68	MMT	300	1.0	2.0	900	80
2009/01/20	+392	0.33–0.87	VLT	300	1.3	1.7	3600	50

^aThe spectra for +32, +58, +122 & +392d are based on data obtained from the ESO Science Archive Facility. The spectra for +96 & +162d are taken from Milisavljevic et al. (2010).

^bWith reference to the epoch of explosion JD 2454462.17.

^cNOT : ALFOSC on 2.6-m Nordic Optical Telescope (NOT), la Palma; NTT : EMMI on 3.6-m New Technology Telescope, la Silla; VLT : FORS2 on 8.2-m ESO-VLT-UT1 telescope, Paranal Observatory; MMT : Blue Channel spectrograph on 6.5-m MMT at Mt. Hopkins;

^dSignal-to-noise-ratio at $0.6\mu\text{m}$.

resents the position of $\text{H}\alpha$. We have marked few nebular lines like $\text{H}\alpha$, $[\text{O II}] \lambda 3727$, which belong to the associated star forming region and is appeared starting from the early spectrum. The absence of Silicon (Si) feature at $6315\text{\AA}$ as well as the absence of H lines indicate that the spectra of SN 2007uy resemble those of stripped-envelope SNe. The lines are highly blended with each other and for most of the lines, the P-Cygni profiles are affected by ‘line-blanketing’. The metallic lines like $\text{Fe II } \lambda 4401$, $\text{Mg II } \lambda 4481$ as well as $[\text{Fe II}] \lambda 5536$, $\text{Na I D } \lambda \lambda 5890, 5896$ and $\text{Ca II } \lambda \lambda 8496, 8542$ are prominent from the early epochs. In the +17d spectrum, the features between $4700\text{\AA}$ and $5000\text{\AA}$ are plausibly attributed by the blend of $\text{He I } \lambda 4731$ along with Fe II multiplates. The lines of O, Mg and Ca start to appear nearly $\sim 30\text{d}$ after the explosion. The spectra evolved faster than other Type Ibc SNe and except $[\text{O I}] \lambda \lambda 6300, 6364$; $[\text{Ca II}]$ and $[\text{O II}]$, all other features almost faded out by +162d. This is in contrast with normal Type Ibc SNe, where emission line profiles remain prominent even beyond +250d (see for e.g., Cox 2000; Modjaz et al. 2008; Milisavljevic et al. 2010; Valenti et al. 2011).

The spectral lines $[\text{O I}] \lambda 5577$ and $[\text{O I}] \lambda \lambda 6300, 6364$ show highly blueshifted emission profiles in the early phase spectra, before eventually moving to their respective rest wavelengths at the +92d phase and onwards. The high blueshift of the spectral features is a signature of aspheric explosion. The blueshift of the spectral lines at early epochs was also observed in other Type I events, especially in the case of Type Ia events (Maeda et al. 2010a,b, 2011).

In figure 3 we compare the +17d spectrum of SN 2007uy with the spectra of Type Ia, Ia-pec, Ibc and Ic events, taken at similar phases. The blueshift in the bluer part of the spectrum is clearly visible; and, the spectrum of SN 2007uy is highly similar to that of Type Ia and Ia-pec events (Maeda et al. 2010a,b; 2011). The characteristics that discriminate SN 2007uy from Type Ia events are: the presence of He I lines as well as strong $\text{H}\alpha$ which indicates that the progenitor is associated with a star forming region, a common site for Type Ibc events. Moreover SN 2007uy is radio luminous (§2.2), which is the result of interaction of the SN shock with the circumburst medium. Type Ia SNe progenitors are not surrounded by a dense circumstellar medium and hence




Figure 3. Spectroscopic comparison of SN 2007uy with other Type I events near maxima. The colour version of the figure is available in the online journal.

are very rarely observable in the radio in the early epochs (Chomiuk et al. 2012; Weiler et al. 2002).

3.1 Evolution of some spectral lines: a message regarding the aspheric explosion

Figure 4 presents the spectral evolution of $\text{Mg I } \lambda 4571$, $[\text{Fe II}] \lambda 5536$, $\text{Na I D } \lambda \lambda 5890, 5896$, $[\text{O I}] \lambda \lambda 6300, 6364$ and $\text{O I } \lambda 7774$, which are commonly found in Type Ibc events and in Type IIP SNe. During the early epochs most of the features are highly blueshifted with respect to their rest wavelengths / velocities.

The early and late phase blue-shifting of the spectral lines can arise for different reasons, e.g., (i) contamination due to other emission lines (Elmhamdi et al. 2003), (ii) a torus or elongated ejecta with a sufficiently opaque in-




Figure 4. Temporal evolution of some spectral lines of SN 2007uy. The zero velocity shown with dotted line in each panel refers the rest wavelength of corresponding elements as mentioned at the top of the panels. The flux scale is relative. To make the features prominent, the intensity of each feature has been normalized by the flux of the zeroth velocity of the corresponding feature, though all the flux related measurements have been done using the normalized spectra shown in figure 2.

ner portion (Chugai 1992; Wang & Hu 1994) or (iii) due to dust formation (Danziger et al. 1989; Lucy et al. 1989; Elmhamdi et al. 2003). For the formation of dust which can block the red wings of the spectral lines, the presence of a cold SN shell is required, whereas newly formed dust grains can only increase the continuum part (as discussed in §5.2). With a Monte-Carlo simulation Taubenberger et al. (2009) showed that, due to the residual opacity of the core ejecta, the emission lines can be blueshifted only about by 30 – 40 Å, which is much less than the blueshifts ($\sim 70\text{--}100\text{Å}$) ob-

served for SN 2007uy. Thus, the third possibility can be ruled out for young ($\lesssim 100\text{d}$) hot SN ejecta.

Asphericity in the SN explosion, with a torus or disk-like distribution of different elements perpendicular to a semi bipolar jet, could also be an explanation for the aspheric signature of the lines viz, double-horn emission profile of [O I] $\lambda\lambda 6300, 6364$ (Maeda et al. 2008; Modjaz et al. 2008 and references therein). The residual opacity in the core of elongated ejecta may also be responsible for the observed blueshift. Optically thick inner ejecta could obstruct the light from the rear side of the SN, creating a flux

deficit in the redshifted part of the emission lines (Chugai 1992; Wang & Hu 1994). It is worth mentioning that symmetric double-peaked profiles can also be explained by the doublet nature of [O I] $\lambda\lambda 6300, 6364$ seen under optically thick conditions leading to an intensity ratio close to one (Milisavljevic et al. 2010). Here we discuss the spectral evolution of each feature.

The Mg I] $\lambda 4571$ (the first panel from the left of figure 4) is seen clearly in +35d spectrum as a broad (FWHM $\sim 185\text{\AA}$), highly blue-shifted ($\sim 71\text{\AA}$) emission peak that corresponds to $\sim 4660\text{ km s}^{-1}$ blueshifted velocity of the emitting region projected onto the line of sight. The ‘double-horn’ feature marked with A at the extreme blue end, is also seen in +17d spectrum. The Mg I] feature starts to move towards its rest wavelength after its first appearance at +32d and remains as a broad emission peak (FWHM $\sim 205\text{\AA}$) until +122d with the emission peak at zero velocity. This broad feature almost disappears in the +162d spectrum and remains as a weak feature marked as B in the deep nebular spectrum obtained at +392d. The feature B is actually blended with a relatively blue-shifted broad emission feature C. After deblending these two features, we marked the feature C as Ba II $\lambda 4554$ (Cox 2000).

The appearance of the extremely blueshifted (velocity $\sim -17500\text{ km s}^{-1}$) emission peak of the blended lines of [O I] $\lambda 5577$, [Fe II] $\lambda 5536$ and [Co II] $\lambda 5526$ at +17d can be seen clearly (the second panel from the left of figure 4). A similar blueshifted emission was initially identified as [O I] $\lambda 5577$ for SN 1993J (Spyromilio 1994; Filippenko et al. 1994; Wang & Hu 1994) and for SN 2007uy (Milisavljevic et al. 2010), although the study of Houck & Fransson (1996) showed that this emission is a blend of [O I], [Fe II] and [Co II]. Over time this feature developed and started to shift toward its rest wavelength from +122d onwards. This confirms the prediction of Houck & Fransson (1996) and establishes this emission line as a blend. Similar to Mg I], [Fe II] $\lambda 5536$ also showed a symmetric emission profile at around zero velocity in late epoch ($\gtrsim 122\text{d}$) spectra. Interestingly from +122d (labeled 5 in the second panel, from left, in figure 4) the intensity of this blended line dropped drastically. The blueshifted emission peak of [O I] $\lambda 5577$ in the late time evolution of the SN ejecta is commonly observed in Type Ibc SNe, e.g. 2004ao, 2006T, 2008D and 2008bo (Milisavljevic et al. 2010), but in none of the cases was a corresponding redshifted component was found.

The Na I D feature (the third panel from the left of figure 4) is more or less a perfect P-Cygni profile, nonetheless it also shows a highly blueshifted (velocity $\sim 7700\text{ km s}^{-1}$) emission peak, possibly blended with the He I $\lambda 5876$ emission feature. The corresponding He I absorption dip is marked as D at +17d. The velocity drops down to about 1933 km s^{-1} at +32d and the feature becomes almost like a P-Cygni profile from +58d. The Na I D feature is absent in +392d spectrum and the He I $\lambda 5876$ emission feature of the underlying star forming region is clearly visible at its rest wavelength. A careful inspection shows the existence of a tiny emission peak E and an absorption dip F at ‘zero velocity’, starting from the initial epoch (+32d). We speculate that E arises from of He I, while F is a footprint of intervening Na I D absorption due to the host galaxy. There are two tiny absorption features around E – one is at the right F and other at the left, about 2000 km s^{-1} blueshifted with

respect to F that is equal to the redshift of the host. Hence the possibility of Na I D impressions due to host (feature F) and Milky Way (blueshifted feature) can not be ruled out.

The [O I] $\lambda\lambda 6300, 6364$ is a frequently studied emission line in the context of the asymmetric nature of Type Ibc explosions. Different geometries of oxygen ejecta have been proposed as the origin of the observed line profiles of this doublet. A flat-topped profile is produced by a radially expanding spherical shell of oxygen gas; a parabolic profile is produced by a filled uniform sphere; and the double horned profile is due to the presence of a cylindrical ring or torus like structure, that expands in the equatorial plane along the line of sight, where the bulk of the emitting gas is located at the projected expansion velocity ($\pm v$) of the torus and a jet-like ejection of matter is along the direction perpendicular to the line of sight. The ‘ $\pm$ ’ sign represents the red and blueshifted projected velocity of the torus respectively (Maeda et al. 2008; Modjaz et al. 2008 and references therein). Alternatively it can be interpreted as the doublet nature of [O I] $\lambda\lambda 6300, 6364$, seen in an optically thick environment (Milisavljevic et al. 2010). In the fourth panel of figure 4 we have shown its evolution. In +32d spectrum a tiny emission feature G is appeared at a blueshifted velocity $\sim 5839\text{ km s}^{-1}$. Apparently it seems that by +58d, this feature moves to a new position G2 along with the appearance of a second peak H at the bluer end. However, due to lack of spectrum between +32d and +58d, this dynamics can not be supported strongly. The velocity of H with respect to rest position of [O I] $\lambda 6300$ is $\sim 7732\text{ km s}^{-1}$. This corresponds to a wavelength separation of about $163\text{\AA}$. Undoubtedly G2 represents the rest wavelength of [O I] $\lambda 6300$ and it has been found to be preserved during follow-up observations. G2 may also be an evolved stage of G1, a very tiny feature just above the noise level in +32d spectrum. The feature G at $6205\text{\AA}$ may be due to a blend of Fe II and Ca II multiplates along with N II, and Ti II, whereas since G1 starts to appear from early epochs, it is possibly due to a shallow spherical distribution of [O I] at the outer ejecta. At +96d the separation between H and G1 is reduced to about $64\text{\AA}$, but in the +122d, +162d and +392d spectra the emission peak profile became almost symmetrical around ‘zero velocity’. This indicates an uniform spherical distribution of [O I] $\lambda\lambda 6300, 6364$ from +122d onward. Since G1 and H do not appear simultaneously, their internal separation is varying with time and the intensity ratio between the blue and red wings of the double horn feature does not tend toward 3:1 with time; we rule out the possibility of appearance of the doublet nature of [O I] $\lambda\lambda 6300, 6364$ in the form of a double horn profile, at least for this particular case. Since these features start to appear from a very early phase, the jet and disk/torus-like scenario, which is invoked to explain late time ($\gtrsim 200$ days) [O I] features, may not be applicable to explain this particular spectral evolution. Similarly the ‘Flat-top’ scenario is also not applicable.

Unlike blended [O I] $\lambda 5577$ and [O I] $\lambda\lambda 6300, 6364$, the O I $\lambda 7774$ feature, shown in the right most panel of figure 4, developed as a symmetrical flat-topped peak, centered around ‘zero velocity’. Although the present spectra do not cover the emergence and development of the O I $\lambda 7774$ line explicitly, they give a clear indication regarding the development of this particular line at around +96d from a completely featureless part of the spectrum. At +122d it be-




Figure 5. Temporal evolution of velocities of SiII and HeI spectral features in SN 2007uy.

comes prominent, though it disappears at +162d. The flat-topped peak is a clear indication of a shell-like structure of the line emitting region and a rapid dilution of the line emitting region is possibly responsible for the sudden decrease of line intensity. At +392d, O I $\lambda 7774$ disappears completely, rather we notice the impression of [Ar III] $\lambda 7751$ at its rest wavelength, about 1000 km s^{-1} blueshifted with respect to O I $\lambda 7774$ in velocity domain.

From the above discussion it seems that the velocity profiles of different line forming regions are not identical for SN 2007uy and hence probably the distribution of ^{56}Ni does not influence the distribution of different material inside the ejecta of SN 2007uy, otherwise the evolution of all material would be of similar. The progenitors of Type Ib SNe have mainly He and O rich shells (Shigeyama et al. 1990). From the geometry of line profiles (Modjaz et al. 2008), we propose that O I $\lambda 7774$ line forming region was distributed inside the ejecta in the form of a shell whereas Na I D and Mg ions were concentrated toward the inner portion of the ejecta with a roughly spherical distribution. Evolution of the blended [O II] $\lambda 5577$ and [O II] $\lambda 6300$ lines are especially interesting. The projected velocity of H show a power law profile with temporal decay index -1.6 where the decay indices of Mg I], Na I D and [O II] $\lambda 5577$ blend (rest wavelength $\sim 5540 \text{ \AA}$) are respectively -1.5 , -1.6 and -1.3 . This confirms that the regions which are forming the lines [O II] $\lambda 5577$ blend, feature H, Mg I] and Na I D are ‘attached’ with each other. Interestingly, the [O II] $\lambda 5577$ blend shows as a very tiny feature (FWHM $\sim 41 \text{ \AA}$) at +32d, before the emergence of feature H and became prominent (FWHM $\sim 63 - 68 \text{ \AA}$) in the +58d and +96d spectra during the appearance of feature H, before again shrinking to a tiny feature with FWHM $\sim 34 \text{ \AA}$ at +122d when G2 and H merged together.

By +122d H meets G2 (or both settle down to a constant

velocity) and starts to move in a spherically symmetric manner and the ejecta becomes sufficiently cool. Our photometric calculation shows that temperature of this ejecta should asymptotically reach below 5000K (see §5.2). In a simulation on forbidden lines in astrophysical-jets Hartigan et al. (2004) showed that below 5000K the strength of [O I] $\lambda 5577$ lessens in comparison to that of [O I] $\lambda 6300$ ($\lesssim 0.1$), even with a moderate electron number density (see figure 9 of Hartigan et al. 2004). Here we speculate that this could be the reason for the disappearance of [O I] $\lambda 5577$ line in between +96d and +122d.

The spectral evolution at the locations of the most prominent He I lines is displayed in figure 5. He I $\lambda\lambda$ 6678, 7065 and 7281 are the strongest and, at the same time, most isolated He lines in the optical spectrum of a SN and are thus often used to distinguish between a SN Ib and Ic. In contrast, He I $\lambda 5876$ is often blended with Na I D (figure 4), a line that also appears in SNe Ic, and He I $\lambda 4471$ is in a region of the spectrum with other overlapping transitions, including Fe lines. Our spectrum at maximum light (+17d) does not show any trace of He in any reasonable velocities. He I $\lambda 7065$ appears 15 days later at a velocity of $\sim 12000 \text{ km s}^{-1}$ (although the line is blended with the telluric B-band). In the same spectrum we do not see any other credible He feature. The same holds for the spectrum at +58d, where He I $\lambda 7065$ has decelerated to 10000 km s^{-1} . He I $\lambda\lambda$ 6678, 7281 finally appear in the +96d spectrum in a consistent velocity with He I $\lambda 7065$ ($\sim 7000 \text{ km s}^{-1}$). Figure 5 also shows the region around $6000 \text{ \AA}$. This region is dominated by an absorption feature that is usually attributed to Si II $\lambda 6355$ in most Type I SNe. Adopting this identification, this line decelerates from $\sim 15000 \text{ km s}^{-1}$ at +17d to $\sim 10000 \text{ km s}^{-1}$ at +96d. What is striking, however, is the appearance of this feature in our second spectrum (+32d) where it has become very strong and broad. It is obvious that this absorption cannot be caused by Si only (it extends to negative velocities) but it must be a blend with more features. Attributing this to He I $\lambda 6678$ is of course a possibility. However, this line would have to appear at very high velocities in order to appear blended with the Si line (in most SNe Ib, these two features are clearly distinguishable). In addition, the line would have to be very strong and that appears to be in conflict both with the velocity and strength of the only other He I line detected in this spectrum (He I $\lambda 7065$) and with the sudden disappearance of this feature in the next spectrum. If we attribute the (now weak) feature on the right of Si II $\lambda 6355$ to He I $\lambda 6678$ in the +58d spectrum, this would have to be at $\sim 5000 \text{ km s}^{-1}$, i.e., a velocity inconsistent with the one inferred by He I $\lambda 7065$ at the same epoch. In general, we note that the appearance and disappearance of this deep trough at $6000 \text{ \AA}$ is rather unusual in comparison to normal Type Ib events.

4 DISTANCE AND EXTINCTION TOWARD SN 2007uy

To determine the bolometric light curve and physical properties of the transient, a correct estimation of the distance and extinction is essential.

The host NGC 2770 is a well studied star forming galaxy

with nine redshift independent distance estimations⁷. Out of them, eight measurements used the ‘Tully Fisher’ technique and for one IRAS photometry was used. The measured values have a range between 26.1 Mpc and 36.0 Mpc, that corresponds to a weighted mean of 29.97 ± 3.48 Mpc. The Hubble flow distance of the host⁸, after correction for Virgo infall, is estimated as 29.3 ± 2.1 Mpc. Combining the above measurements, we adopt the weighted mean distance of 29.5 ± 1.8 Mpc, which corresponds to a distance modulus of ~ 32.4 mag.

The Galactic reddening along the line of sight of SN 2007uy as derived from the $100\mu\text{m}$ all sky dust extinction map (Schlegel et al. 1998) is $E(B - V) = 0.022 \pm 0.0004$ mag. However, measuring the total line-of-sight extinction towards SN 2007uy is non-trivial. SN 2007uy is associated with a large number (about 150) of $\text{H}\alpha$ emitters (Gorosabel et al. 2011) which are expected to be star forming HII regions and presumably few of them can make the transient highly extinguished. The host is also a highly inclined spiral galaxy (see Table 1), where the dusts are normally distributed along the edges of the young and old stellar discs (Popescu et al. 2011 and references therein) and hence expected to produce a large extinction toward the SN due to the foreground material. In fact, it has already been pointed out by Drout et al. (2011), with sufficient number of Type Ib events, that by and large the more extincted SNe reside in more inclined host galaxies. For e.g., SN 2007D is the most extinguished object which is hosted by a highly inclined ($\sim 70^\circ$) galaxy; however, there are exceptions like SN 2004ge.

NGC 2770 hosted three SNe 1999eh, 2007uy and 2008D. The isophotal diameter of NGC 2770 is $d_{25} = 3.567'$. This corresponds to an apparent radius $r_{25} = 1.734'$ of the host. On the other hand since NGC 2770 is a spiral galaxy, assuming the star forming regions and SNe sites are distributed on the plane of the disk, the measured values of the deprojected distances of SNe 2007uy, 2008D and 1999eh from the centre of the host are roughly $0.707'$ ($\equiv 5.3\text{kpc}$), $1.210'$ ($\equiv 9.1\text{kpc}$) and $1.262'$ ($\equiv 9.5\text{kpc}$) respectively. This implies $r_{2007uy} \sim 0.408r_{25}$, $r_{2008D} \sim 0.698r_{25}$ and $r_{1999eh} \sim 0.728r_{25}$. Thus all the SNe happened within half-light radius of the host. Moreover according to Thöne et al. (2009) the metallicities of these SNe sites are also comparable with each other (8.36 ± 0.1 , 8.53 and 8.37 dex respectively for SN 2008D, SN 2007uy, SN 1999eh). So we can expect that the environments are similar and the host extinctions should have nearly equal values.

From the extinction values of the well studied nearby Type Ibc events, Drout et al. (2011) proposed an empirical relation to calculate the host reddening for Type Ibc events using the values of observed $V - R$ colour of the transient at +10d after V and R band maxima. According to this formalism, the magnitudes of the host reddening, $E(B - V)$ for SN 2007uy measured w.r.t the V and R band maxima are respectively 0.79 ± 0.18 mag and 0.82 ± 0.21 mag.

The colour excess of the host can be measured by means of Balmer decrement, usually from $\text{H}\alpha/\text{H}\beta$ ratio (Osterbrock 1989) or from the equivalent width of the non-contaminated and non-saturated NaID absorption feature (Barbon et al. 1990; Richmond et al. 1994; Elias-Rosa et al. 2007; Poznanski et al. 2012).

By measuring the Balmer decrement, Thöne et al. (2009) determined the $E(B - V) \sim 1.4$ mag⁹ toward SN 2007uy. This corresponds to total visual extinction ~ 4.34 mag considering the ratio of total-to-selective extinction (R_V) of 3.1. We note that with this value of visual extinction, the peak absolute magnitude of SN 2007uy would be ~ -21 mag. This is roughly 3.4 mag brighter than the average peak magnitude of Type Ibc events (Drout et al. 2011) and comparable with over-luminous events, though neither the photometric nor the spectroscopic evolution of this event matches with those of luminous SNe. It is needless to say that measurement of the $\text{H}\alpha/\text{H}\beta$ ratio is critically dependent on the correct flux calibration and prescriptions for deriving atmospheric and foreground extinction. The broad spectral lines of a bright SN is also a probable dominating source of error in determination of the ratios of underlying lines. We perform the same exercise with +392d spectrum and calculate the reddening from Balmer decrement (see e.g. Domínguez et al. 2013) using the expression

$$E(B - V) = 1.97 \times \log_{10}[(\text{H}\alpha/\text{H}\beta)_{\text{obs}}/2.86] .$$

The derived value of reddening $E(B - V)$, is 0.62 ± 0.06 mag.

On the other hand, for SN 2007uy we found a prominent impression, F, of the host NaID , though the two components are not resolved in the moderate resolution spectra. Using our four high S/N spectra (+32d, +58d, +96d and +122d) where the unresolved NaID feature was prominent, we calculate its mean equivalent width (EW) of $0.66 \pm 0.04\text{\AA}$. By using the expression of Poznanski et al. (2012) for the unresolved NaID feature we derive the value of reddening to be $E(B - V) = 0.08^{+0.03}_{-0.02}$ mag. We note that this value is similar to the line of sight reddening towards the face-on or less inclined galaxies (Stritzinger et al. 2009; Sahu et al. 2009; Roy et al. 2011a) and too low for a transient hosted by a highly inclined galaxy having a substantial foreground intervening medium. This value is comparable to the lower limit of reddening (0.063) that can be inferred from NaID absorption feature in low resolution spectra (Elias-Rosa et al. 2007). However, in the same contribution it was also pointed out that there is no clear empirical relation between $E(B - V)$ and EW , at least in the regime of low-resolution spectroscopy. We also notice a prominent impression E of $\text{HeI } \lambda 5876$ in the spectra of SN 2007uy, that is due to associated star forming region. This may be a prime source of contamination for the NaID .

To avoid these problems, we measure the line ratio between $\text{NII } \lambda 6583$ and $\text{H}\alpha$ and adopt an indirect method to measure the $E(B - V)$ toward SN 2007uy. $\text{NII } \lambda 6583/\text{H}\alpha$ ratio is actually a good estimator for Oxygen abundance

⁷ <http://www.ned.ipac.caltech.edu/>

⁸ The cosmological model with $H_0 = 70 \text{ km s}^{-1} \text{ Mpc}^{-1}$, $\Omega_m = 0.3$ and $\Omega_\Lambda = 0.7$ is assumed throughout the paper and the uncertainty corresponds to a local cosmic thermal velocity of 208 km s^{-1} (Terry et al. 2002).

⁹ This is important to mention that Thöne et al. (2009) did not calculate the $E(B - V)$ exactly at SN position, rather at the locations which are roughly $\pm 2''$ separations from SN position. This corresponds to more than 350pc in physical size.

in extragalactic H II regions (Pettini & Pagel 2004). After noticing a strong anti-correlation between N II $\lambda 6583$ /H α ratio and H α equivalent width, whereas strong correlations among [O II], H α , H β and H γ fluxes in a sample of about 74 star forming blue galaxies, Kong et al. (2002) proposed an empirical relation between $E(B - V)$ and N II $\lambda 6583$ /H α . They found that since N II $\lambda 6583$ and H α are two nearby lines, the strength ratio will be less affected due to galactic extinction while providing an approximate tracer of $E(B - V)$ for highly reddened ($E(B - V) \gtrsim 0.1$ mag) extragalactic objects. Note, though, this is inefficient for low reddened objects. The +392d spectrum, which has the least SN contribution and mostly contains the emission lines due to H α knots, has been used to calculate $E(B - V)$. The measured value of $E(B - V)$ by this relation is $0.52^{+0.20}_{-0.14}$ mag. This is similar to the reddening along highly reddened SNe 2004ge and 2007D and also close to the mean reddening ($E(B - V) \sim 0.65$ mag) along SN 2008D (Maund et al. 2009).

For this work, we adopt the weighted average of four measurements, obtained from the empirical relations proposed by Kong et al. (2002); Drout et al. (2011) and ratio of Balmer decrement. This implies that the host reddening toward SN 2007uy is 0.63 ± 0.15 mag. Since this is an order of magnitude higher than the galactic extinction, we anticipate that this is the total reddening along the line of sight of SN 2007uy. This corresponds to a total visual extinction (A_V) of 1.9 ± 0.5 mag.

5 LIGHT CURVE EVOLUTION: NUV, OPTICAL AND NIR

The calibrated light curves of SN 2007uy are presented in figure 6. The data cadences of ground based Optical and NIR observations are $N(U, B, V, R, I, J, H, K) = (7, 19, 23, 20, 22, 8, 8, 8)$, while that for spaced based NUV and Optical observations are $N(uvw2, uvm2, uvw1, u, b, v) = (14, 8, 18, 19, 28, 30)$ respectively. Since the host redshift is small, no ‘K-correction’ is applied. From the peak, the SN light is mainly powered by the radioactive decay of ^{56}Ni and ^{56}Co . The energy of γ -rays and positrons emitted by radioactive decay processes is fully absorbed by the SN ejecta and re-emitted as blackbody radiation. Hence in early phases Type I events can be approximated as ‘blackbody supernova’ (Arnett 1982). The emergence of a young Type I SN with its blackbody nature is clearly noticeable in all bands. At a few phases, the SN was not detected in some of the *Swift*/UVOT bands for which the corresponding upper limits are mentioned. In the ground-based observations, the maximum sampling of SN light was done in the V passband. Similarly, for the space-based *Swift*/UVOT observations the maximum sampling was done in v passband. Since the central wavelength and bandwidth of the V passband (5448Å and 840Å respectively) is fairly similar to that of the v passband (5468Å and 769Å respectively), we have treated the combined data of the V and v bands as a single visual-data set for further analysis of photometric data and likewise the B and b -band data sets have been combined as a single blue-band data set. In contrast, a significant deviation between the U and u passband light curves, starting from early epochs ($\gtrsim 20$ days), is probably due to a considerable

difference between their responses (for the U band: central wavelength 3663Å bandwidth 650Å for the u band: central wavelength 3465Å bandwidth 785Å respectively).

The evolution of the SN light curve is composed of three distinct phases – (i) shock breakout phase (initial few days), (ii) photospheric phase (within first 40–50 days) and (iii) nebular phase (beyond 50 days after explosion). For this particular event we will discuss the evolution of photospheric and nebular phase light curves.

5.1 Evolution during photospheric phase

The temporal variation of the visual light roughly resembles the temporal variation of the total UVOIR (NUV+Optical+NIR) light of the SN. The maxima of visual light also roughly corresponds to the epoch of maximum of the UVOIR bolometric light of the transient. The rise time of SN 2007uy in the visual band is 19.4 day which is determined after estimating the maximum through third order polynomial fitting on the early part ($\lesssim 40$ days) of the visual light curve. This is comparable with the rise time derived for engine driven energetic, Type Ic SN 1998bw (16.14 ± 0.08 days) and optically-normal Type Ib SN 2008D (18.82 ± 0.24 days), whereas, it is longer than that of Type Ic SN 2002ap (10.12 ± 0.20 days) and engine driven energetic SN 2006aj (10.37 ± 0.14 days). The measured rise time of SN 2007uy in U band is 15.13 ± 0.4 days, whereas in I band it is 23.09 ± 1.44 days. This shows that, similar to other Ibc events, SN 2007uy also evolved faster and peaked first at lower wavelengths and then at higher wavelengths.

The peak width ($\Delta d_{0.25}$), defined as the width of the light curve, when the SN has faded by 0.25 mag from its maximum brightness, is also calculated for SN 2007uy. For the visual band it is roughly 11 days. This is comparable to the V band peak width of SN 2006aj ($\Delta d_{0.25} \sim 11.1$ days) and SN 2002ap ($\Delta d_{0.25} \sim 10.6$ days), but less than engine driven SN 1998bw ($\Delta d_{0.25} \sim 13.3$ days) and normal Type Ib event SN 2008D ($\Delta d_{0.25} \sim 18.04$ days).

The post maxima decay rates of SN 2007uy in the V and R bands are respectively 0.058 ± 0.002 mag.d $^{-1}$ and 0.042 ± 0.002 mag.d $^{-1}$, which corresponds to post maxima decay parameters of (Δm_{15}) of 0.87 ± 0.003 and 0.63 ± 0.03 mag respectively. Here Δm_{15} quantifies the decay in magnitude in 15 days after maxima. For the B and U bands, the value of Δm_{15} are even higher, around 1.2 and 2.1 mag respectively. This is in agreement with the prediction of models on SN evolution (Woosley et al. 1999b). The V and R band decay rates are comparable with the decay rates of Type Ibc events studied by Drout et al. (2011). On the other hand, the spectroscopic study (see §3) showed convincingly that the event neither belongs to the Type Ic nor to the class Ic-BL.

Adopting the distance modulus of SN 2007uy ~ 32.4 mag and total reddening along the line of sight $E(B - V) \sim 0.63$ mag (see §4), we have computed absolute magnitudes in the $BVRI$ passbands to compare it with other Type Ibc events (figure 7). For relatively redshifted objects, like SN 2006aj, the cosmological time dilation factor has been taken care into account to correct the corresponding time scale. In all bands SN 2007uy was intrinsically brighter than the normal Type Ib event SN 2008D and Type Ic event SN 2002ap. Though in the R and I pass-




Figure 6. The light curves of SN 2007uy in NIR *JHK*, Optical *UBVRI* and *uvw2*, *uvm2*, *uvw1&u*, *b*, *v* NUV-Optical UVOT bands. The colour version of the figure is available in the online journal.

bands it was a little dimmer than the GRB/XRF associated events SNe 2006aj and 1998bw, but its peak absolute *R* band magnitude -18.5 ± 0.16 is about one mag brighter than the mean value (-17.6 ± 0.6) derived for well observed Type Ibc events (Drout et al. 2011). In the visual and blue bands its peak magnitude is comparable with energetic events associated with GRB/XRFs. Hence, SN 2007uy is optically more luminous than normal Type Ib events and comparable to the engine driven energetic SNe.

SN 2007uy was luminous in the NUV. It was detectable in the *Swift*/*uvw2* and *uvm2* images until +24d, while in the *uvw1* and *u* filter systems it was detected until +44d. In contrast, SN 2008D which was more luminous in the X-ray

and radio and happened in the same galaxy with almost similar environment (Thöne et al. 2009), was seldom detected in the *Swift* NUV bands. As we have noticed that SN 2007uy is luminous in higher optical frequencies and it is comparable to engine driven energetic explosion like SN 2006aj, it is worth to compare the NUV flux of this event with that of engine driven SNe. Fortunately, *Swift* has a good coverage of XRF060218/SN 2006aj in all NUV bands during the phases comparable to that of SN 2007uy. Figure 8 shows a comparison between the NUV and *u* absolute magnitudes of SN 2007uy with that of SN 2006aj. The UVOT data for SN 2006aj have been taken from Šimon et al. (2010). SN 2006aj seems to be dimmer than SN 2007uy. The nature of




Figure 7. Absolute light curves of SN 2007uy. Comparison with other Type Ib and Ic events.




Figure 8. NUV and u band absolute light curves of SN 2007uy. Comparison with Type Ic event SN 2006aj. Data for SN 2006aj were presented by Šimon et al. (2010). The colour version of the figure is available in the online journal.

the evolving supernova is clearly visible in all absolute light curves of SN 2007uy. In the case of SN 2006aj, the nature of an evolving blackbody supernova is reflected in the *u* and marginally in the *uvw1* light curves, whereas photometry of *uvm2* and *uvw2* images show a very shallow linear decay of SN light in the NUV range. The post maxima decay rates in these two bands, determined through a linear fit to these data sets, are respectively 0.06 mag d^{-1} and 0.04 mag d^{-1} .

If all optical photons are generated through blackbody radiation under similar conditions of the ejecta, then post maxima decay rates in these two bands should be much higher than these values (assuming the SN will be optically thick during first 20 days); it was also predicted by Woosley et al. (1999a) through modeling of SN 1998bw spectra in the spectral range between the *U* and *I* passbands. For SN 2006aj, the measured post-maxima decay rate in the *I* band is $\sim 0.05 \text{ mag d}^{-1}$ while in the *B* and *uvw1* bands these are as high as $\sim 0.10 \text{ mag d}^{-1}$ and 0.12 mag d^{-1} respectively. Hence the decay rate in NUV bands should be higher. This discrepancy was also noted by Šimon et al. (2010) and was explained as an effect of blackbody radiation emission (Panagia 2003) while the UV emitting area shrinks with time during the evolution of the transient.

In contrast, blackbody fitting to the early SN 2007uy data (figure 9) shows that envelope of this SN evolved more or less like a blackbody. Here we have plotted the flux, normalized with respect to the *V* band flux, to nullify the effect due to time evolution of the angular diameter of the transient. The effect due to dilution of SN envelope is also neglected as a first approximation¹⁰. The temperature of each blackbody is mentioned inside the parenthesis, with a temperature-scale 1000 K. The one sigma uncertainties are measured by applying the ‘Monte Carlo’ method. The initial temperature of the photosphere is $\sim 8000^\circ \text{ K}$. During the next 6 days the temperature slightly increased to $\sim 8100^\circ \text{ K}$ and then started to decrease with time through a power law with temporal decay index ~ -0.51 , which is comparable with the theoretical prediction (-0.5) for a ‘blackbody supernova’ by Arnett (1982). The initial increment in temperature, followed by a power law decay with time (decay index ~ -0.12) was also observed in the Type Ic SN 2002ap (Vinkó et al. 2004). Thus we speculate that unlike XRF060218/SN 2006aj, SN 2007uy evolved roughly as a ‘blackbody supernova’ from the very initial epoch.

5.2 Evolution during nebular phase

Normally after +50 to +60 days, the envelopes of most Type I events start to become optically thin, and hence the light curves start to flatten. In the *B*, *V*, *R* and *I* bands the decay rate of SN 2007uy is roughly $0.017 - 0.018 \text{ mag d}^{-1}$, whereas in *U* band it is $\sim 0.03 \text{ mag d}^{-1}$. This decay rate is similar to the nebular decay rates of other Type Ibc events and slightly higher than the decay rate of the $^{56}\text{Co} \rightarrow ^{56}\text{Fe}$ nuclear transition, which is typically $0.0098 \text{ mag d}^{-1}$. It is worth mentioning that in Type II SNe, mainly in normal Type IIP events, having a huge H shell in the outer surface, the nebular decay rates are comparable with that of ^{56}Co

¹⁰ It is worth noting that the measurement of the true Temperature of the SN envelope along with the measurement of its distance require a proper estimation of dilution factor through hydrodynamical modeling (Kirshner & Kwan 1974; Eastman et al. 1996). This can only provide a colour temperature for the system along with a true estimation of distance independent of cosmology. The technique was properly applied by Hamuy et al. (2001); Leonard et al. (2002) and others for Type IIP events. In the present work we provide a rough estimate of the overall temperature of the SN photosphere and adopted the value of distance obtained above (§4).




Figure 9. Temporal evolution of observed SED of SN 2007uy. The temperature of each blackbody is mentioned inside the parenthesis, with a temperature-scale 1000K. The one sigma uncertainties are measured by applying the ‘Monte Carlo’ method. The colour version of the figure is available in the online journal.

(Roy et al. 2011a,b and references therein). On the other hand, for low luminosity Type IIP events like SN 2005cs, the decay rate during nebular phase ($\sim 0.0046 \text{ mag d}^{-1}$ in V band during the period 140–320 days) is significantly lower than that of ^{56}Co (Pastorello et al. 2009). This was also observed by Elmhamdi et al. (2003) in the normal Type IIP event SN 1999em during its initial ($\sim$ one month) nebular phase. Utrobin (2007) explained this as an effect due to radiation generated inside the inner warmed ejecta, that propagates through the cool, optically thin outer region and makes the nebular light relatively stable. The relatively high decay in very late epoch ($t > 300$ days) during the evolution of SN 2005cs was explained as resulting either due to (i) dust formation in the SN ejecta, (ii) a lower efficiency of γ -ray trapping due to the decreased density of the ejecta, or due to (iii) cooling of inner ejecta (Pastorello et al. 2009).

The optical and NIR nebular light curves of SN 2007uy show sharper decay ($\sim 0.02\text{--}0.03 \text{ mag.d}^{-1}$) than the canonical decay of radioactive ^{56}Co . A similar decay rate was also noticed for SN 2002ap (Pandey et al. 2003; Vinkó et al. 2004), which was explained as a leakage of γ -photons from the transparent and less massive SN atmosphere. Dust formation may also be another cause. With increasing time, the newly formed small dust grains in the ejecta can progressively reduce the fluxes of SN in all bands. Though there is a possibility for the creation of dust-grains of size $\lesssim 0.01 \mu\text{m}$ within the first 100–300 days in Type I SNe (Smith et al. 2008; Nozawa et al. 2011), these hot grains ($T \sim 1600^\circ \text{K}$) would cause an IR excess in the SN continuum spectra. Moreover, dust formation should manifest itself by creating a blueshift in the line, increasing with time. Neither excess in the continuum, nor the temporal increment in the blueshifts of the spectral lines were observed for SN 2007uy (rather it

decreases with time; see §3). In fact beyond +50d SN 2007uy started to become bluer (see §6), which does not support dust formation during the period of observation. Thus the possibility of dust formation at the outer shell of the ejecta in early epoch is also unable to explain this deviation.

Here we propose that leakage of γ -photons is the principal cause for rapid decay in light curve of SN 2007uy, like other stripped-envelope supernovae. We also admit that this simultaneously depends on the amount of ejected mass (M_{ej}) and how fast it becomes diluted (optically thin). If E_k is the kinetic energy of the explosion then the decay rate of the nebular phase is roughly proportional to $M_{ej}/E_k^{1/2}$ (Valenti et al. 2008). So, explosions with higher ejecta mass and kinetic energy or lower mass with low kinetic energy should show a similar trend in nebular decay. For e.g., the very late time (> 150 days after explosion) evolution of SNe 1998bw and 2006aj also show an almost identical decay rate in the V , R , I bands of $\sim 0.02 \text{ mag.d}^{-1}$ and higher than the decay rate of ^{56}Co (Patat et al. 2001; Misra et al. 2011). The first one is an explosion with higher mass of ejecta and kinetic energy, while the second shows a lower ejecta mass with smaller amount of kinetic energy.

6 EVOLUTION OF COLOUR AND BOLOMETRIC LIGHT

The reddening-corrected colour evolution of SN 2007uy is presented in figure 10. For comparison, the colour curves of the well-studied SNe 1998bw (Galama et al. 1998), 2002ap (Foley et al. 2003), 2006aj (Ferrero et al. 2006) and 2008D (Soderberg et al. 2008; Roy 2013) have been plotted. The evolution of intrinsic colour of SN 2007uy is similar to that of other events, however during early phases ($\lesssim +35\text{d}$) the SN seems to be bluer than the other SNe, especially in $(V-I)_0$ colour. This is a manifestation of enhanced line emission due to $\text{He I } \lambda 5876$ and $[\text{O I}] \lambda 5577$ in comparison to the higher wavelength part of the spectrum, which is also confirmed from +17d spectrum. Like other events, SN 2007uy also shows a transition from blue to red during its initial evolution ($\lesssim +40\text{d}$) and turned bluish during its nebular evolution ($\gtrsim +50\text{d}$). This is probably due to transition of ejecta from optically thick to optically thin state, which in turn initiates the escape of high energy optical photons in a large amount from the SN ejecta.

The bolometric light curve is crucial for the calculation of the synthesized radioactive ^{56}Ni , the kinetic energy of the explosion, the amount of mass that was expelled during the explosion and hence to understand the nature of the progenitor. The quasi-bolometric UVOIR light curve has been computed from UV, optical and NIR data for the event that is presented in figure 11. Bolometric fluxes have been computed in those epochs for which visual magnitudes are available. The extinction-corrected magnitudes are first converted into fluxes using zero points given by Bessell et al. (1998) and then the total flux in the UVOIR bands is obtained after a linear interpolation and integration between 0.203 and $2.19 \mu\text{m}$. The JHK contribution during the photospheric phase is calculated from our data, whereas beyond +80d, in the nebular phase, due to lack of UV and NIR data, we are not able to make any direct measurement. For many Type Ia supernovae it has been observed that the NUV




Figure 10. Color evolution of SN 2007uy. Comparison with other Type Ibc events

and NIR contribution at the nebular phase is roughly 20% (Contardo et al. 2000; Leloudas et al. 2009). However, for Type Ic SN 2002ap, Tomita et al. (2006) found a relatively higher contribution ($\sim 25\text{--}30\%$) from NIR light during comparable epochs. So, for present analysis, beyond +80d, we increase the bolometric flux by 35% to estimate the maximum possible contribution from NUV and NIR light and construct the UVOIR quasi-bolometric light curve. Similarly, before the peak, there are few epochs for which we do not have any *RI* and *JHK* data. To calculate the magnitudes at those epochs, a blackbody spectrum has been fitted to construct the SED for each day (see figure 9) and synthetic magnitudes have been calculated from the SED.

The peak UVOIR bolometric luminosity of SN 2007uy is about $6.4 \times 10^{42} \text{ erg s}^{-1}$ which is comparable to that of the engine-driven energetic SN 1998bw ($\sim 7.5 \times 10^{42} \text{ ergs}^{-1}$) and SN 2006aj ($\sim 5.9 \times 10^{42} \text{ ergs}^{-1}$), and much higher than the broad-lined Type Ic event SN 2002ap ($\sim 2.1 \times 10^{42} \text{ ergs}^{-1}$) and the optically-normal Type Ib event SN 2008D ($\sim 2.0 \times 10^{42} \text{ ergs}^{-1}$). The peak luminosity of Type I SNe is roughly proportional to the ejected amount of ^{56}Ni because the contributions from shock heating and CSM interaction are small. Since the maximum light of SN 2007uy is almost similar to that of SNe 2006aj and 1998bw, hence the amount of synthesized radioactive ^{56}Ni nuclei should be comparable with these engine driven events. On the other hand, the decay rate in nebular phase is comparable with decay of ^{56}Co to ^{56}Fe , though there is a slight deviation as mentioned in §5.2.

The bolometric light of a CCSN can comprise light from different components, such as the explosion of a single massive progenitor, the emergence of a magnetar, the interaction of ejecta with the circumstellar medium (CSM), or possibly due to a combination of these. The signature of SN shock-interaction with its CSM can be seen through radio observations (see §7).




Figure 11. The UVOIR bolometric light curve of SN 2007uy. Comparison with other Type Ibc events. The colour version of the figure is available in the online journal.




Figure 12. The modeled UVOIR bolometric light curve of SN 2007uy. The black dots are the observed data points. The blue solid line represents the model. The colour version is available in online journal.

7 PHYSICAL PARAMETERS OF THE EXPLOSION

To determine the physical parameters of the exploding star, modeling of the UVOIR bolometric light is essential. In addition to that, to quantify the interaction with circumstellar medium, modeling of the radio data is also required.

7.1 Modeling of optical light curve

To derive the physical parameters, we have followed the methodology of Valenti et al. (2008), i.e. the early and nebular phase have been modeled separately.

For $t \lesssim 30$ days past explosion, the ejecta is presumed to be optically thick, having a spherically symmetric, homologous expansion and all the radioactive ^{56}Ni is located in the

centre. The model also assumes that the radius of the expanding shell is much higher than the radius of the progenitor, the optical opacity $\kappa_{opt} \sim 0.06 \text{ cm}^2 \text{ g}^{-1}$ remains constant through out its evolution (Maeda et al. 2003), and the total radioactive energy of $^{56}\text{Ni} \rightarrow ^{56}\text{Co} \rightarrow ^{56}\text{Fe}$ is responsible for the heating of the ejecta and the diffusion approximation is valid for the photons (Arnett 1982; Valenti et al. 2008; Benetti et al. 2011; Pignata et al. 2011; Olivares E. et al. 2012). In the nebular phase (beyond $\gtrsim 60$ days), the ejecta become optically thin and the emitted luminosity is powered by three sources – instantaneous energy of the γ -rays generated during the ^{56}Co decay, energy of the γ -rays coming from electron-positron annihilation and due to the kinetic energy of the positrons (Sutherland & Wheeler 1984; Cappellaro et al. 1997). The low and high γ -ray trapping, respectively at early and late phases, have been invoked by dividing the ejecta into two components – a high-density inner region and a low-density outer region (Maeda et al. 2003). The outer region dominates the total emission mechanism at early epochs, i.e. in the optically thick regime, and emission from the inner region, with higher γ -ray opacity, dominates the nebular phase.

Optical modeling, as mentioned above, returns the values of four free parameters used to fit the bolometric light curve of SN 2007uy. These are: the total mass of the ejected material (M_{ej}), the total mass of ^{56}Ni produced in the envelope (M_{Ni}), the mass fraction of the inner ejecta component (f_M) and the fraction of kinetic energy contributed by the inner component (f_E). To break the degeneracy between kinetic energy and ejected mass, the expression for velocity at peak luminosity has been used (Arnett 1982; also see the equation 2 of Valenti et al. 2008 and equation 3 of Olivares E. et al. 2012).

Velocities of the ejecta are measured from the blue shifts of the absorption dips of the P-Cygni profiles, but due to the heavy line blanketing identification of photospheric lines is not trivial and also the possibility to get the undisturbed lines is much less. Our first spectrum, taken near maximum, contains the He I $\lambda 5876$ line, with a considerable blue-shifted absorption dip. The measured velocity corresponding to this blue shift is roughly 15200 km s^{-1} . Assuming the He shells to be marginally detached from the photosphere, we can constrain the upper limit of the photospheric velocity (v_{ph}) within the above value.

The fitted model is presented in figure 12. The parameters obtained after modeling the observed bolometric light of this event are as follows : $M_{ej} \sim 4.4 M_\odot$, $M_{Ni} \sim 0.3 M_\odot$, $f_M \sim 9\%$ and $f_E \sim 0.02\%$. Since the peak velocity and total ejected mass are related to the Kinetic energy (E_K) of the entire ejecta by the expression:

$$E_K = [(5/6) M_{ej} (1 - f_M) v_{ph}^2] / (1 - f_E),$$

the kinetic energy of the ejecta turned out to be $E_K \sim 15 \times 10^{51}$ ergs.

7.2 Modeling of radio data

The radio light curve can be explained in terms of the evolution of the ejecta, which further determines the shape of the spectrum and evolution of its peak with time (Sari et al. 1998; Wijers & Galama 1999). It can also be explained by Synchrotron Self Absorption (SSA) with high brightness




Figure 13. The modeled radio light curve of SN 2007uy. The upper panel shows the variation of radio signal with time. Open circles are the data obtained from van der Horst et al. (2011), whereas the filled circles are VLA data acquired for the present work. The solid curves represent the best fit model. The lower panel shows the variation of spectral index with time.

temperature and/or by a Free Free Absorption (FFA) from the ambient medium, which produces an exponential cut-off below the peak. A decrease in absorption with time, resulting a smooth, rapid turn-on, first at higher frequencies and later at lower frequencies. For each frequency, the peak flux corresponds to optical depth ~ 1 and a power-law decline in the light curve can be seen. Finally, the spectral index approaches a constant negative value, which corresponds to non-thermal radiation in an optically thin medium (Weiler et al. 1986, 1990; Chevalier 1998) and Weiler et al. (1986) proposed a methodology for parameterization of SN radio light curve.

For the present work, we have parameterized the radio light curves of SN 2007uy to calculate the mass loss rate. The Upper panel of figure 13 presents the results of radio observations in six different frequencies. The early time data is too sparse to determine the spectral index (α) for the early part of the light curve, during the optically thick regime. Radio follow-up of this event has already been reported by van der Horst et al. (2011).

To model the light curves we have also included the literature data along with this new data set. The modeled parameters are reported in Table 8. The notation used for the parameters is as defined by Weiler et al. (2011). The details for the modeling can be found in Weiler et al. (1986, 1990, 2011).

Out of the six frequencies of the data set, only three (8.4, 4.8 and 2.3 GHz) have been used to model the radio data through simultaneous fitting, while for other frequencies (1.28, 0.610 and 0.325 GHz) the expected theoretical light curves have been plotted along with observational data. Our best fitting model gives the value of overall reduced-

Table 8. The parameters derived from the fitting of radio data

Parameters	Value
K_1	1500.75
α	-1.1435
β	-1.734
K_2	91.38
t_0	-3.59 days
δ	2.41
K_3	5.408×10^5
K_4	0.0036
δ'	-3.43
K_5	1630.03
δ''	-4.60
K_6	~ 1.0008
δ'''	-4.20

$\chi^2 = 7.14$. This is relatively higher than the value (4.42) mentioned by Weiler et al. (2011) for the well-studied Type Ibc radio SN 1994I. This higher value is presumably a reflection of a lack of data due to less coverage of SN 2007uy in radio band. The lower panel of figure 13 presents the temporal evolution of α . The epochs for which we have an observation in a given frequency, the probable flux for other frequencies was determined by interpolation through polynomial fitting of the observed light curve. The thick curve shows that initially the source was extremely optically-thick with high +ve value of spectral index which turned to a constant -ve value ~ -1 asymptotically, that clearly marks the transition from a optically thick to thin state. A lower value (~ 2.0) of α than 2.5 in the optically thick regime shows FFA is also responsible for the initial radio flux along with the SSA. Although, the coefficient $K_6 = 1.00085$, which is associated with FFA is also sufficiently less than the coefficient $K_5 = 1630.031$ responsible for SSA, the temporal evolution of both processes are almost similar ($\delta'' = -4.602$ and $\delta''' = -4.206$ respectively). Thus the initial emission is mostly due to SSA in optically thick ejecta, though a mild contribution due to FFA can not be ruled out. This result is consistent with the conclusion drawn by van der Horst et al. (2011).

The modeling of the three frequencies shows that the radio flux of the source is moderate ($K_1 = 1500.75$) with a temporal decay index (β) roughly $= -1.73$ while the overall spectral index (α) is ~ -1.14 , comparable with the optically thin condition. It predicts that the object was discovered about (t_0) $= -3.6$ days after the explosion, which is consistent with the optical follow-ups. Following Chevalier (1982a,b), we assume that the temporal decrement (δ) of the opacity of a uniform circumstellar medium (CSM) is roughly dependent on α and β by the relation $\delta = \alpha - \beta - 3$. A sufficiently lower value of K_2 ($= 91.38$), than K_3 ($= 5.408 \times 10^5$) implies that the absorption processes in CSM are mainly due to its clumpy/structured form than its uniformity. On the other hand a relatively higher value of δ' ($= -3.43$) than δ ($= -2.41$) implies that the effect of clumpiness will not sustain for a longer time in comparison to the absorption due to uniform CSM. So, eventually the CSM starts to behave like a circumburst medium having a statistically small number of clumps. The lower value of the term K_4 ($= 3.6 \times 10^{-3}$) de-

mands that the radio absorption due to FFA in distant H II regions, which are in between the SN and us is quite low. In fact this is one order less than other events likes SNe 1978K, 1994I and 1998bw, but higher than SN 1993J (Weiler et al. 2002, 2011). This is in contrast with the large extinction toward SN 2007uy, derived from optical data (§4).

Finally, we find the mass loss rate during the pre-SN phase using radio data, adopting the prescription of Weiler et al. (2001). Assuming 50% of the CSM along the line of sight is clumpy and the size of a clump is roughly 1/3 of the dimension of the expanding SN shock, then the volume filling factor by the blast wave is $\phi \sim 0.22$. The effective optical depth will be given by $\langle \tau_{eff}^{0.5} \rangle \sim 0.204$ (using equations 5 and 16 of Weiler et al. 2001). Assuming the wind velocity of the progenitor $\sim 10 \text{ km s}^{-1}$ (the least possible value if the progenitor is a red supergiant) and initial velocity (v_s) and electron temperature (T_s) of the shock are respectively of the order of $20,000 \text{ km s}^{-1}$ and $20,000 \text{ K}$, we can quantify the mass loss rate by:

$$\dot{M} = 6.79 \times 10^{-7} (w_{wind}/10 \text{ km s}^{-1}) \cdot (v_s/20,000 \text{ km s}^{-1})^{1.5} \cdot (T_s/20,000 \text{ K})^{0.68} \text{ M}_{\odot} \text{ yr}^{-1}$$

Since the photospheric velocity and temperature, measured from spectroscopy and photometry at around optical peak are roughly $15,200 \text{ km s}^{-1}$ and $8,000 \text{ K}^{11}$ respectively, then the mass loss rate should be $\dot{M} \gtrsim 2.41 \times 10^{-7} (w_{wind}/10 \text{ km s}^{-1}) \text{ M}_{\odot} \text{ yr}^{-1}$. This value is comparable with the pre-SN mass loss rate for other Type Ibc events derived from radio data (Weiler et al. 2011). Observations and theoretical studies predict that the progenitors of Type Ibc are WR stars, having extremely high wind velocity $\gtrsim 10^3 \text{ km s}^{-1}$ (Yoon et al. 2012; Willis 1996). Then the mass loss rate of the progenitor of SN 2007uy during its WR phase was roughly $\gtrsim 2.41 \times 10^{-5} \text{ M}_{\odot} \text{ yr}^{-1}$. Certainly this result is consistent with other Type Ibc events (Weiler et al. 2011).

8 CONCLUSIONS

Since the detection of the progenitors of Type Ibc events in pre-SN images is a technically challenging task, rigorous observational follow-up along with theoretical interpretation is necessary to understand these catastrophic explosions in a better way.

In this article, we have presented an extensive follow-up campaign of a normal Type Ib event SN 2007uy, using the data acquired in a wide range of wavelength – from NUV to radio bands along with optical spectroscopy. The radio modeling predicts that the event was discovered within 4 days after the explosion, while the early spectral features constrain the time difference between explosion and detection within 7 days. We have adopted the former value and assumed that the SN was discovered within 4 days after the explosion. The main results are summarized below.

¹¹ This is important to mention that this is the colour temperature of the photosphere and hence does not correspond to the electron temperature of the source but provides an approximate estimation of the source temperature.

The extinction ($A_V = 1.9 \pm 0.5$ mag) along the line of sight of SN 2007uy is mainly dominated by the host galaxy. The highly inclined host and the presence of numerous H α emitters along the line of sight make the transient highly extinguished.

The evolution of SN 2007uy was like a canonical ‘black-body supernova’. The UV component is probably not due to shock breakout, rather it is a manifestation of radioactively heated, relatively less massive ejecta. As the ejecta become progressively optically thin, the optical photons start to escape rapidly, causing a steepening in the late time optical light curve.

The spectral evolution of SN 2007uy is faster than other Type Ibc events. Most of the lines get diluted quickly in comparison to other archetypal events. Early evolution of most of the lines is highly aspheric, though their natures are different from each other. This demonstrates that there is no dependency of asphericity on the distribution of radioactive ^{56}Ni inside the ejecta, at least for this particular case. We propose that SN 2007uy is an aspheric explosion, which in due course of time attained a large-scale spherical symmetry.

The nature of colour evolution of SN 2007uy is similar to other stripped-envelope SNe though this transient is intrinsically more bluish than other Type Ibc events. SN 2007uy is one of the luminous Type Ib events in respect to its UVOIR bolometric flux. The peak flux is comparable with the engine driven SNe 1998bw and 2006aj.

The basic parameters of the explosion have been derived from the optical and radio data modeling. The UVOIR optical data has been modeled assuming the event as a ‘point explosion’ after adopting the formalism of Arnett (1982), along with the modifications introduced by Valenti et al. (2008). Optical modeling predicts that about $4.4M_\odot$ was ejected with an explosion energy $\sim 15 \times 10^{51}$ erg and roughly $0.3M_\odot$ was produced during this explosion. The radio data shows a SSA dominated light curve evolution of SN 2007uy, though the contribution of FFA during the early phase can not be ruled out. The line of sight is probably dominated by relatively cool gas and dust making the optical extinction high, but due to lack of free-electron content, absorption of radio waves through FFA is less. It is one order less than that for other radio SNe. The pre-SN mass loss rate obtained from the radio data is $\dot{M} \gtrsim 2.03 \times 10^{-7}$ ($w_{\text{wind}}/10\text{km s}^{-1}$) $M_\odot \text{ yr}^{-1}$. This value is consistent with the results obtained for other Type Ibc radio SNe, though an order of magnitude lower than typical value of mass-loss rate derived from X-ray. Assuming a WR star as a progenitor with wind velocity $\gtrsim 10^3 \text{ km s}^{-1}$, we can constrain the lower limit of pre-SN mass loss rate at $\dot{M} \gtrsim 2.03 \times 10^{-5} M_\odot \text{ yr}^{-1}$.

ACKNOWLEDGMENTS

We thank all the observers at ARIES who provided their valuable time and support for the observations of this event. We are thankful to the observing staffs of NOT, UKIRT, NTT, MMT and VLT for their kind cooperation in observation of SN 2007uy. This work is based on the data obtained from the ESO Science Archive Facility. For this research work the VLA radio data has been used. The VLA is operated by the National Radio Astronomy Observatory, a facility of the National Science Foundation operated under co-

operative agreement by Associated Universities, Inc. DARK is funded by the DNRf. Giorgos Leloudas is supported by the Swedish Research Council through grant No. 623-2011-7117. Rupak Roy is thankful to the COSPAR fellowship program under which, he got the opportunity to visit DARK and initiated the entire project. S. B. Pandey and Rupak Roy acknowledge DST-RFBR grants INT/RFBR/P-25 (2008-2010) and INT/RFBR/P-100 (2011-2013) for the present work. This research has made use of *Swift*/UVOT data obtained through the High Energy Astrophysics Science Archive Research Center (HEASARC) Online Service, provided by the NASA/Goddard Space Flight Center. The authors would also like to thank the anonymous referee for the comments and suggestions which helped in improvement of this manuscript.

REFERENCES

- Arnett W. D., 1982, ApJ, 253, 785
- Barbon R., Benetti S., Rosino L., Cappellaro E., Turatto M., 1990, A&A, 237, 79
- Benetti S. et al., 2011, MNRAS, 411, 2726
- Bessell M. S., Castelli F., Plez B., 1998, A&A, 333, 231
- Blondin S., Calkins M., 2008, Central Bureau Electronic Telegrams, 1191, 2
- Burrows A., 2013, Rev. Mod. Phys., 85, 245
- Cappellaro E., Mazzali P. A., Benetti S., Danziger I. J., Turatto M., della Valle M., Patat F., 1997, A&A, 328, 203
- Chevalier R. A., 1982a, ApJL, 259, L85
- Chevalier R. A., 1982b, ApJ, 259, 302
- Chevalier R. A., 1998, ApJ, 499, 810
- Chomiuk L. et al., 2012, ApJ, 750, 164
- Chugai N. N., 1992, Soviet Astronomy Letters, 18, 168
- Clocchiatti A., Wheeler J. C., 1997, ApJ, 491, 375
- Contardo G., Leibundgut B., Vacca W. D., 2000, A&A, 359, 876
- Cox A. N., 2000, Allen’s astrophysical quantities
- Danziger I. J., Gouffes C., Bouchet P., Lucy L. B., 1989, IAU Circ., 4746, 1
- Domínguez A. et al., 2013, ApJ, 763, 145
- Drout M. R. et al., 2011, ApJ, 741, 97
- Eastman R. G., Schmidt B. P., Kirshner R., 1996, ApJ, 466, 911
- Eldridge J. J., Fraser M., Smartt S. J., Maund J. R., Crockett R. M., 2013, ArXiv e-prints
- Elias-Rosa N., Beckman J. E., Benetti S., Cappellaro E., Turatto M., 2007, in Supernovae: lights in the darkness
- Elmhamdi A. et al., 2003, MNRAS, 338, 939
- Ferrero P. et al., 2006, A&A, 457, 857
- Filippenko A. V., Matheson T., Barth A. J., 1994, AJ, 108, 2220
- Foley R. J. et al., 2003, PASP, 115, 1220
- Galama T. J. et al., 1998, Nature, 395, 670
- Gorosabel J. et al., 2011, Advances in Space Research, 47, 1421
- Hamuy M. et al., 2001, ApJ, 558, 615
- Hartigan P., Edwards S., Pierson R., 2004, ApJ, 609, 261
- Heger A., Fryer C. L., Woosley S. E., Langer N., Hartmann D. H., 2003, ApJ, 591, 288
- Hirst P., Casali M., Adamson A., Ives D., Kerr T., 2006,

- in Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series, Vol. 6269, Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series
- Höflich P., Khokhlov A., Wang L., 2001, in American Institute of Physics Conference Series, Vol. 586, 20th Texas Symposium on relativistic astrophysics, Wheeler J. C., Martel H., eds., pp. 459–471
- Horne K., 1986, *PASP*, 98, 609
- Houck J. C., Fransson C., 1996, *ApJ*, 456, 811
- Kirshner R. P., Kwan J., 1974, *ApJ*, 193, 27
- Kong X., Cheng F. Z., Weiss A., Charlot S., 2002, *A&A*, 396, 503
- Kumar B., Sagar R., Rautela B. S., Srivastava J. B., Srivastava R. K., 2000, *Bulletin of the Astronomical Society of India*, 28, 675
- Landolt A. U., 1992, *AJ*, 104, 340
- Leloudas G. et al., 2009, *A&A*, 505, 265
- Leonard D. C. et al., 2002, *AJ*, 124, 2490
- Lucy L. B., Danziger I. J., Gouffes C., Bouchet P., 1989, in *Lecture Notes in Physics*, Berlin Springer Verlag, Vol. 350, IAU Colloq. 120: Structure and Dynamics of the Interstellar Medium, Tenorio-Tagle G., Moles M., Melnick J., eds., p. 164
- Maeda K. et al., 2010a, *Nature*, 466, 82
- Maeda K. et al., 2008, *Science*, 319, 1220
- Maeda K. et al., 2011, *MNRAS*, 413, 3075
- Maeda K., Mazzali P. A., Deng J., Nomoto K., Yoshii Y., Tomita H., Kobayashi Y., 2003, *ApJ*, 593, 931
- Maeda K., Taubenberger S., Sollerman J., Mazzali P. A., Leloudas G., Nomoto K., Motohara K., 2010b, *ApJ*, 708, 1703
- Malesani D. et al., 2009, *ApJL*, 692, L84
- Maund J. R., Smartt S. J., 2005, *MNRAS*, 360, 288
- Maund J. R., Smartt S. J., Schweizer F., 2005, *ApJL*, 630, L33
- Maund J. R., Wheeler J. C., Baade D., Patat F., Höflich P., Wang L., Clocchiatti A., 2009, *ApJ*, 705, 1139
- Maund J. R., Wheeler J. C., Patat F., Baade D., Wang L., Höflich P., 2007, *MNRAS*, 381, 201
- Mazzali P. A. et al., 2005, *Science*, 308, 1284
- Milisavljevic D., Fesen R. A., Gerardy C. L., Kirshner R. P., Challis P., 2010, *ApJ*, 709, 1343
- Misra K., Fruchter A. S., Nugent P., 2011, in *American Institute of Physics Conference Series*, Vol. 1358, American Institute of Physics Conference Series, McEnery J. E., Racusin J. L., Gehrels N., eds., pp. 299–302
- Modjaz M., Kirshner R. P., Blondin S., Challis P., Matheson T., 2008, *ApJL*, 687, L9
- Nakano S., Kadota K., Itagaki K., Corelli P., 2008, *Central Bureau Electronic Telegrams*, 1191, 1
- Nozawa T., Maeda K., Kozasa T., Tanaka M., Nomoto K., Umeda H., 2011, *ApJ*, 736, 45
- Olivares E. F. et al., 2012, *A&A*, 539, A76
- Osterbrock D. E., 1989, *Astrophysics of gaseous nebulae and active galactic nuclei*
- Panagia N., 2003, in *Lecture Notes in Physics*, Berlin Springer Verlag, Vol. 598, Supernovae and Gamma-Ray Bursters, Weiler K., ed., pp. 113–144
- Pandey S. B., Anupama G. C., Sagar R., Bhattacharya D., Sahu D. K., Pandey J. C., 2003, *MNRAS*, 340, 375
- Pastorello A. et al., 2009, *MNRAS*, 394, 2266
- Patat F. et al., 2001, *ApJ*, 555, 900
- Pettini M., Pagel B. E. J., 2004, *MNRAS*, 348, L59
- Pignata G. et al., 2011, *ApJ*, 728, 14
- Poole T. S. et al., 2008, *MNRAS*, 383, 627
- Pooley D., Soderberg A., 2008, *The Astronomer's Telegram*, 1368, 1
- Popescu C. C., Tuffs R. J., Dopita M. A., Fischera J., Ky-lafis N. D., Madore B. F., 2011, *A&A*, 527, A109
- Poznanski D., Prochaska J. X., Bloom J. S., 2012, *ArXiv e-prints*
- Pumo M. L. et al., 2009, *ApJL*, 705, L138
- Richmond M. W., Treffers R. R., Filippenko A. V., Paik Y., Leibundgut B., Schulman E., Cox C. V., 1994, *AJ*, 107, 1022
- Roy R., 2013, PhD thesis, Department of Physics, Kumaun University
- Roy R. et al., 2011a, *ApJ*, 736, 76
- Roy R. et al., 2011b, *MNRAS*, 414, 167
- Sahu D. K., Tanaka M., Anupama G. C., Gurugubelli U. K., Nomoto K., 2009, *ApJ*, 697, 676
- Sari R., Piran T., Narayan R., 1998, *ApJL*, 497, L17
- Schlegel D. J., Finkbeiner D. P., Davis M., 1998, *ApJ*, 500, 525
- Shigeyama T., Nomoto K., Tsujimoto T., Hashimoto M.-A., 1990, *ApJL*, 361, L23
- Smith N., Foley R. J., Filippenko A. V., 2008, *ApJ*, 680, 568
- Smith N., Li W., Filippenko A. V., Chornock R., 2011, *MNRAS*, 412, 1522
- Soderberg A., 2008, *The Astronomer's Telegram*, 1350, 1
- Soderberg A. M. et al., 2008, *Nature*, 453, 469
- Spyromilio J., 1994, *MNRAS*, 266, L61
- Stetson P. B., 1987, *PASP*, 99, 191
- Stritzinger M. et al., 2009, *ApJ*, 696, 713
- Sutherland P. G., Wheeler J. C., 1984, *ApJ*, 280, 282
- Taubenberger S. et al., 2009, *MNRAS*, 397, 677
- Terry J. N., Paturel G., Ekholm T., 2002, *A&A*, 393, 57
- Thöne C. C., Michałowski M. J., Leloudas G., Cox N. L. J., Fynbo J. P. U., Sollerman J., Hjorth J., Vreeswijk P. M., 2009, *ApJ*, 698, 1307
- Tomita H. et al., 2006, *ApJ*, 644, 400
- Utrobin V. P., 2007, *A&A*, 461, 233
- Šimon V., Pizzichini G., Hudec R., 2010, *A&A*, 523, A56
- Valenti S. et al., 2008, *MNRAS*, 383, 1485
- Valenti S. et al., 2011, *MNRAS*, 416, 3138
- van der Horst A. J. et al., 2011, *ApJ*, 726, 99
- van Dokkum P. G., 2001, *PASP*, 113, 1420
- Vinkó J. et al., 2004, *A&A*, 427, 453
- Wang L., Howell D. A., Höflich P., Wheeler J. C., 2001, *ApJ*, 550, 1030
- Wang L., Hu J., 1994, *Nature*, 369, 380
- Wang L., Wheeler J. C., 2008, *ARA&A*, 46, 433
- Weiler K. W., Panagia N., Montes M. J., 2001, *ApJ*, 562, 670
- Weiler K. W., Panagia N., Montes M. J., Sramek R. A., 2002, *ARA&A*, 40, 387
- Weiler K. W., Panagia N., Sramek R. A., 1990, *ApJ*, 364, 611
- Weiler K. W., Panagia N., Stockdale C., Rupen M., Sramek R. A., Williams C. L., 2011, *ApJ*, 740, 79
- Weiler K. W., Sramek R. A., Panagia N., van der Hulst J. M., Salvati M., 1986, *ApJ*, 301, 790

- Wijers R. A. M. J., Galama T. J., 1999, ApJ, 523, 177
Willis A. J., 1996, Ap&SS, 237, 145
Woosley S. E., Eastman R. G., Schmidt B. P., 1999a, ApJ, 516, 788
Woosley S. E., MacFadyen A. I., Heger A., 1999b, ArXiv Astrophysics e-prints
Yoon S.-C., Gräfener G., Vink J. S., Kozyreva A., Izzard R. G., 2012, A&A, 544, L11